arXiv:1602.06160v3 [math.NT] 23 Nov 2016

# A Weighted Divisor Problem[☆]

Lirui Jia

*School of Mathematical Sciences, Zhejiang University, Hangzhou 310027, People's Republic of China*

Wenguang Zhai

*Department of Mathematics, China University of Mining and Thechnology, Beijing 100083, People's Republic of China*

**Abstract**

We study a weighted divisor function $\sum_{mn\leq x}{}' \cos(2\pi m\theta_1)\sin(2\pi n\theta_2)$, where $\theta_i (0<\theta_i<1)$ is a rational number. By connecting it with the divisor problem with congruence conditions, we establish an upper bound, mean-value, mean-square and some power-moments.

*Keywords:* Weighted divisor problem, power-moment, upper bound
*2010 MSC:* 11N37, 11P21

## 1. Introduction and main results

### *1.1. Introduction*

Let $d(n)=\sum_{n=n_1n_2}1$ denote the divisor function, and $D(x)=\sum_{n\leq x}{}' d(n)=\sum_{n_1n_2\leq x}{}' 1$ be the summatory function, where the prime $'$ on the summation sign indicates that if $x$ is an integer, then only $\frac{1}{2}d(x)$ or $\frac{1}{2}$ for $n_1n_2=x$ is counted. In 1849, Dirichlet first proved that

$$D(x)=x\log x+(2\gamma-1)x+O(\sqrt{x}),\forall x\geq 1,$$

where $\gamma$ is the Euler constant.

Let

$$\Delta(x)=D(x)-x\log x-(2\gamma-1)x-\frac{1}{4}$$

[☆]The work of Lirui Jia is supported by the National Natural Science Foundation of China (Grant No. 11571303). Wenguang Zhai is supported by the National Key Basic Research Program of China (Grant No. 2013CB834201), the National Natural Science Foundation of China (Grant No. 11171344).

*Email addresses:* jialirui@126.com (Lirui Jia), zhaiwg@hotmail.com (Wenguang Zhai)

be the error term in the asymptotic formula for $D(x)$. Dirichlet's divisor problem consists of determining the smallest $\alpha$, for which $\Delta(x) \ll x^{\alpha+\varepsilon}$ holds for any $\varepsilon > 0$. Clearly, Dirichlet's result above implies that $\alpha \leq \frac{1}{2}$. Throughout the past more that 160 years, there have been many improvements on this estimate. The best estimate to-date has been given by Huxley[4, 5], and reads

$$\Delta(x) \ll x^{\frac{131}{416}} \log^{\frac{26947}{8320}} x. \tag{1.1}$$

It is widely conjectured that $\alpha = \frac{1}{4}$ is admissible and the best possible.

Since $\Delta(x)$ exhibits considerable fluctuations, one natural way to study the upper bounds is to consider the moments.

Voronoi's work[16] in 1904 showed that

$$\int_1^X \Delta(x)dx = \frac{X}{4} + O(X^{\frac{3}{4}}).$$

Later, in 1922 Cramér[2] proved the mean square formula

$$\int_1^X \Delta(x)^2 dx = cX^{\frac{3}{2}} + O(X^{\frac{5}{4}+\varepsilon}), \quad \forall\ \varepsilon > 0,$$

where $c$ is a positive constant. In 1983, Ivic [6] used the method of large values to prove that

$$\int_1^X |\Delta(x)|^A dx \ll X^{1+\frac{A}{4}+\varepsilon}, \quad \forall\ \varepsilon > 0 \tag{1.2}$$

for each fixed $0 \leq A \leq \frac{35}{4}$. The range of $A$ can be extended to $\frac{262}{27}$ by the estimate (1.1). In 1992, Tsang[14] obtained the asymptotic formula

$$\int_1^X \Delta(x)^k dx = c_k X^{1+\frac{k}{4}} + O(X^{1+\frac{k}{4}-\delta_k}), \quad \text{for } k = 3, 4, \tag{1.3}$$

with positive constants $c_3$, $c_4$, and $\delta_3 = \frac{1}{14}$, $\delta_4 = \frac{1}{23}$. Ivić and Sargos [8] improved the values $\delta_3$, $\delta_4$ to $\delta_3' = \frac{7}{20}$, $\delta_4' = \frac{1}{12}$, respectively. Heath-Brown[3] in 1992 proved that for any integer $k < A$, where $A$ satisfies (1.2), the limit

$$c_k = \lim_{X\to\infty} X^{-1-\frac{k}{4}} \int_1^X \Delta(x)^k dx$$

exists. Then, there followed a series of investigations on explicit asymptotic formulas of the type (1.3) for larger values of $k$. In 2004, Zhai [17] established asymptotic formulas for $3 \leq k \leq 9$.

At the beginning of 20th century, Voronoï[16] proved the remarkable exact formula that

$$\Delta(x) = -\frac{2}{\pi}\sqrt{x}\sum_{n=1}^{\infty}\frac{d(n)}{\sqrt{n}}\big(K_1(4\pi\sqrt{nx}) + \frac{\pi}{2}Y_1(4\pi\sqrt{nx})\big),$$

where $K_1$ is the modified Bessel function of the second kind of order 1, and $Y_1$ is the Bessel function of the second kind of order 1. The series on the right-hand side is boundedly convergent for $x$ lying in each fixed closed interval.

Recently, Berndt et al [1] considered $\sum'_{mn\le x}\cos(2\pi m\theta_1)\sin(2\pi n\theta_2)$, which can be seen as a weighted divisor function and they got an analogue of Voronoi's formula as follows.

Let $J_1$ be the ordinary Bessel function. If $0<\theta_1,\theta_2<1$ and $x>0$, then

$$\begin{aligned}&\sum_{mn\le x}{}' \cos(2\pi m\theta_1)\sin(2\pi n\theta_2)\\ =&-\frac{\cot(\pi\theta_2)}{4}+\frac{\sqrt{x}}{4}\sum_{m=0}^{\infty}\sum_{n=0}^{\infty}\Bigg\{\frac{J_1\big(4\pi\sqrt{(m+\theta_1)(n+\theta_2)x}\big)}{\sqrt{(m+\theta_1)(n+\theta_2)}}\\ &+\frac{J_1\big(4\pi\sqrt{(m+1-\theta_1)(n+\theta_2)x}\big)}{\sqrt{(m+1-\theta_1)(n+\theta_2)}}-\frac{J_1\big(4\pi\sqrt{(m+\theta_1)(n+1-\theta_2)x}\big)}{\sqrt{(m+\theta_1)(n+1-\theta_2)}}\\ &-\frac{J_1\big(4\pi\sqrt{(m+1-\theta_1)(n+1-\theta_2)x}\big)}{\sqrt{(m+1-\theta_1)(n+1-\theta_2)}}\Bigg\}.\end{aligned}$$

In this paper, we will study $\sum'_{mn\le x}\cos(2\pi m\theta_1)\sin(2\pi n\theta_2)$ further with a different approach and prove some results analogous to those for $\Delta(x)$.

NOTATIONS. For a real number $t$, let $[t]$ be the largest integer no greater than $t$, $\{t\}=t-[t]$, $\psi(t)=\{t\}-\frac12$, $\|t\|=\min(\{t\},\,1-\{t\})$, $e(t)=e^{2\pi it}$. $\mathbb{C}$, $\mathbb{R}$, $\mathbb{Z}$, $\mathbb{N}$ denote the set of complex numbers, of real numbers, of integers, and of natural numbers, respectively; $C^r[a,b]$ be the class of functions having a continuous $r$th derivative in $[a,b]$; $f\asymp g$ means that both $f\ll g$ and $f\gg g$ hold. Throughout this paper, $\varepsilon$, $\varepsilon'$ denote sufficiently small positive constants, and $\mathcal{L}$ denotes $\log T$.

*1.2. Main results*

We first state the divisor problem with congruence conditions. A divisor function with congruence conditions is defined by

$$d(n;r_1,q_1,r_2,q_2)=\sum_{\substack{n=n_1n_2\\ n_i\equiv r_i\,(\mathrm{mod}\ q_i)\\ i=1,2}}{}' 1,$$

of which, the summatory function is

$$D(x;r_1,q_1,r_2,q_2)=\sum_{\substack{n_1n_2\le x\\ n_i\equiv r_i\,(\mathrm{mod}\ q_i)\\ i=1,2}}{}' 1.$$

From Richert [11], we can find that for $x\ge q_1q_2$, $1\le r_i\le q_i$ $(i=1,2)$

$$\begin{aligned}&D(x;r_1,q_1,r_2,q_2)\qquad(1.4)\\ =&\frac{x}{q_1q_2}\log\Big(\frac{x}{q_1q_2}\Big)-\bigg(\frac{\Gamma'}{\Gamma}\Big(\frac{r_1}{q_1}\Big)+\frac{\Gamma'}{\Gamma}\Big(\frac{r_2}{q_2}\Big)+1\bigg)\frac{x}{q_1q_2}+\Delta(x;r_1,q_1,r_2,q_2),\end{aligned}$$

where

$$\Delta(x;r_1,q_1,r_2,q_2)=-\sum_{\substack{n_1\le\sqrt{\frac{q_1x}{q_2}}\\ n_1\equiv r_1\,(\mathrm{mod}\,q_1)}}\psi\big(\frac{x}{q_2n_1}-\frac{r_2}{q_2}\big)-\sum_{\substack{n_2\le\sqrt{\frac{q_2x}{q_1}}\\ n_2\equiv r_2\,(\mathrm{mod}\,q_2)}}\psi\big(\frac{x}{q_1n_2}-\frac{r_1}{q_1}\big)+O(1).$$

Denote $S(x;\theta_1,\theta_2)=\sum_{mn\le x}{}'\cos(2\pi m\theta_1)\sin(2\pi n\theta_2)$. Let $\theta_1$, $\theta_2$ be rational numbers satisfying $\theta_i=\frac{a_i}{q_i}$, with $(q_i,a_i)=1$ $(i=1,2)$. Then $S(x;\theta_1,\theta_2)=S(x;\frac{a_1}{q_1},\frac{a_2}{q_2})$. For simplicity, in this paper we consider $S(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2})$.

Our first result is an analogue of (1.2) for $\Delta(x;r_1,q_1,r_2,q_2)$, which would play an important role in the study of higher power moments of $S(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2})$.

**Theorem 1.1.** *Suppose $A\ge 0$ is a fixed real number, $T\gg(q_1q_2)^{\varepsilon}$, then*

$$\int_1^T|\Delta(q_1q_2x;r_1,q_1,r_2,q_2)|^Adx\ll\begin{cases}T^{1+\frac{A}{4}}\mathcal{L}^{4A}, & \text{if } 0\le A\le\frac{262}{27};\\ T^{\frac{131A+154}{416}}\mathcal{L}^{4A+2}, & \text{if } A>\frac{262}{27}.\end{cases}$$

**Remark 1.1.** *Let $A_0>1$ be a constant such that*

$$\int_1^T|\Delta(q_1q_2x;r_1,q_1,r_2,q_2)|^{A_0}dx\ll T^{1+\frac{A_0}{4}+\varepsilon}. \tag{1.5}$$

*Then Theorem 1.1 shows that we can take $A_0=\frac{262}{27}$. The value $\frac{262}{27}$, which is obtained by using* (2.1)*, is closely related to the upper bound of $\Delta(q_1q_2x;r_1,q_1,r_2,q_2)$. It is easy to prove that the conjecture* (2.2) *is equivalent to* (1.5) *holding for any $A_0>0$.*

For $S\big(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2}\big)$, we have the following results.

**Theorem 1.2.** (***Upper bound***) *For $x\ge 1, 1\le a_i\le q_i$ and $(a_i,q_i)=1$ $(i=1,2)$, we have*

$$S\big(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2}\big)\ll q_1q_2x^{\frac{131}{416}}(\log x)^{\frac{26947}{8320}}.$$

**Remark 1.2.** *Conjecture* (2.2) *implies that*

$$S\big(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2}\big)\ll q_1q_2x^{\frac{1}{4}+\varepsilon}.$$

**Theorem 1.3.** (***Mean value***) *For a large number $T$, if $1\le a_i\le q_i$ and $(a_i,q_i)=1$ $(i=1,2)$, then*

$$\int_1^T S\big(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2}\big)dx\ll q_1q_2T^{\frac{3}{4}}.$$

**Theorem 1.4.** (***Mean-square***) *Suppose $T \gg (q_1q_2)^{\varepsilon'}$ is large enough, $q_1 \geq 2$, $q_2 \geq 3$, $1 \leq a_i \leq q_i$ and $(a_i, q_i) = 1$ $(i = 1, 2)$. Then*

$$\int_1^T S^2\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big)dx = \frac{(q_1q_2)^2}{2^6\pi^2} B_2(\frac{a_1}{q_1}, \frac{a_2}{q_2}) \int_1^T x^{\frac{1}{2}}dx + O\big((q_1q_2)^2T^{\frac{5}{4}}\mathcal{L}^3\big),$$

*where the constant*

$$B_2\big(\frac{a_1}{q_1}, \frac{a_2}{q_2}\big) = \sum_{n=1}^{\infty} \frac{\big(\Delta d_2(n; a_1, q_1, a_2, q_2)\big)^2}{n^{\frac{3}{2}}} \asymp 1.$$

**Theorem 1.5.** (***Higher-power moments***) *Let $T \gg (q_1q_2)^{\varepsilon'}$ be a large number, $q_1 \geq 2$, $q_2 \geq 3$, $1 \leq r_i \leq q_i$ and $(a_i, q_i) = 1$ $(i = 1, 2)$. If $A_0 > 9$ satisfies* (1.5)*, then for any fixed integer $3 \leq k < A_0$, we have*

$$\begin{aligned} &\int_1^T S^k\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big)dx \qquad (1.6)\\ =&\frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k} B_k(\frac{a_1}{q_1}, \frac{a_2}{q_2}) \int_1^T x^{\frac{k}{4}}dx + O\big((q_1q_2)^kT^{1+\frac{k}{4}-\frac{A_0-k}{4(A_0-2)s(K_0)}+\varepsilon}\big), \end{aligned}$$

*where*

$$B_k(\frac{a_1}{q_1}, \frac{a_2}{q_2}) = \sum_{\upsilon=1}^{k-1} \cos\Big(\frac{3\pi(k-2\upsilon)}{4}\Big)\binom{k-1}{\upsilon} \sum_{\{n_1,\dots n_k\}\in\mathcal{S}(k,\upsilon)} \prod_{i=1}^{k} \frac{\Delta d_2(n_i; a_1, q_1, a_2, q_2)}{n_i^{\frac{3}{4}}},$$

*with $\Delta d_2(n; a_1, q_1, a_2, q_2)$, $K_0, s(K_0)$ and $\mathcal{S}(k, \upsilon)$ defined in Section 3. Clearly, $B_k \asymp 1$.*

**Remark 1.3.** *From Theorem 1.1 we can get the asymptotic formula* (1.6) *for $k \in \{3, 4, \dots, 9\}$, by taking $A_0 = \frac{262}{27}$. Moreover,* (1.6) *provides the exact form of the main term for $\int_1^T S^k\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big)dx$ with $k \geq 10$, and if the conjecture* (2.2) *were true, then we could get an asymptotic formula for $\int_1^T S^k\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big)dx$ with any $k \geq 3$.*

If $A_0 = \frac{262}{27}$, then $K_0 = 10$, and $s(K_0) = 257$. Hence, we get an immediate corollary of Theorem 1.5.

**Corollary 1.1.** *Let $3 \leq k \leq 9$ be a fixed integer. If $T, a_i$ and $q_i (i = 1, 2)$ satisfying the hypothesis of Theorem 1.5, then*

$$\int_1^T S^k\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big)dx = \frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k} B_k(\frac{a_1}{q_1}, \frac{a_2}{q_2}) \int_1^T x^{\frac{k}{4}}dx + O\big((q_1q_2)^kT^{1+\frac{k}{4}-\frac{262-27k}{213824}+\varepsilon}\big).$$

**Remark 1.4.** *There are similar conclusions for $\sideset{}{'}\sum_{mn\leq x} \sin(2\pi n\theta_1)\sin(2\pi m\theta_2)$ and $\sideset{}{'}\sum_{mn\leq x} \cos(2\pi n\theta_1)\cos(2\pi m\theta_2)$, which we do not present in this paper.*

**Acknowledgement.**The authors deeply thank the referee for carefully reading the manuscript and valuable suggestions.

## 2. Background of divisor problem with congruence conditions

In this section, we review some background of the divisor problem with congruence conditions.

*2.1. Some estimates for* $\Delta(x; r_1, q_1, r_2, q_2)$

It follows from Huxley's estimates[4] that

$$\Delta(x; r_1, q_1, r_2, q_2) \ll \Big(\frac{x}{q_1 q_2}\Big)^{\frac{131}{416}} \Big(\log\Big(\frac{x}{q_1 q_2}\Big)\Big)^{\frac{26947}{8320}} \tag{2.1}$$

uniformly in $1 \le r_1 \le q_1 \le x, 1 \le r_2 \le q_2 \le x$. It is conjectured that

$$\Delta(x; r_1, q_1, r_2, q_2) \ll \Big(\frac{x}{q_1 q_2}\Big)^{\frac{1}{4}+\varepsilon} \tag{2.2}$$

uniformly in $1 \le r_1 \le q_1 \le x, 1 \le r_2 \le q_2 \le x$, $\forall \varepsilon > 0$, which is an analogue of the well-known conjecture that $\Delta(x) \ll x^{\frac{1}{4}+\varepsilon}$.

Müller and Nowak[10] studied the mean value of $\Delta(x; r_1, q_1, r_2, q_2)$. They pointed out

$$\int_1^T \Delta(x; r_1, q_1, r_2, q_2)dx \ll (q_1 q_2)^{-\frac{1}{4}} T^{\frac{3}{4}} \tag{2.3}$$

and

$$\int_1^T \Delta^2(x; r_1, q_1, r_2, q_2)dx = c_2 (q_1 q_2)^{\frac{1}{2}} T^{\frac{3}{2}} + o\big((q_1 q_2)^{\frac{1}{2}} T^{\frac{3}{2}}\big) \tag{2.4}$$

uniformly in $1 \le r_i \le q_i \le T$ $(i = 1, 2)$, if $T$ is a large number, and $c_2$ is a constant.

*2.2. An analogue Voronoi formula for* $\Delta(x; r_1, q_1, r_2, q_2)$

In this subsection we shall prove the following.

**Lemma 2.1.** *Let* $J = [\frac{\mathcal{L}+2\log q_1 q_2 - 4\log \mathcal{L}}{\log 2}]$, $H \ge 2$ *be a parameter to be determined, and* $T^\varepsilon < y \le \min(H^2, (q_1 q_2)^2 T)\mathcal{L}^{-4}$. *Suppose* $\frac{T}{2} \le x \le T$. *Denote*

$$\tau(n, x) = \sum_{n=hl} \cos\left(4\pi\sqrt{nx} - 2\pi\Big(\frac{hr_2}{q_2} + \frac{lr_1}{q_1} + \frac{1}{8}\Big)\right),$$

$$\tau_{12}(n, x; H) = \sum_{\{h,l\}\in E(n;H,J)}{}' \cos\left(4\pi\sqrt{nx} - 2\pi\Big(\frac{hr_2}{q_2} + \frac{lr_1}{q_1} + \frac{1}{8}\Big)\right),$$

$$\tau_{21}(n, x; H) = \sum_{\{h,l\}\in E(n;H,J)}{}' \cos\left(4\pi\sqrt{nx} - 2\pi\Big(\frac{hr_1}{q_1} + \frac{lr_2}{q_2} + \frac{1}{8}\Big)\right),$$

*with* $E(n; H, J)$ *defined in Section 3. Then*

$$\begin{aligned}\Delta(q_1 q_2 x; r_1, q_1, r_2, q_2) = {} & R_0'(x; y) + {R_{12}}'(x; y, H) + {R_{21}}'(x; y, H) \\ & + G_{12}'(x; H) + G_{21}'(x; H) + O\big(\log^3(q_1 q_2 T)\big).\end{aligned} \tag{2.5}$$

*Here*

$$R_0'(x;y)=\frac{x^{\frac14}}{\sqrt{2}\pi}\sum_{n\le y}\frac{\tau(n,x)}{n^{\frac34}},$$

$$R_{12}{}'(x;y,H)=\frac{x^{\frac14}}{\sqrt{2}\pi}\sum_{y<n\le 2^{J+1}H^2}\frac{\tau_{12}(n,x;H)}{n^{\frac34}},$$

$$R_{21}{}'(x;y,H)=\frac{x^{\frac14}}{\sqrt{2}\pi}\sum_{y<n\le 2^{J+1}H^2}\frac{\tau_{21}(n,x;H)}{n^{\frac34}},$$

$$G_{12}'(x;H)=\sum_{\substack{n_1\le q_1\sqrt{T}\\ n_1\equiv r_1 \,(\mathrm{mod}\ q_1)}} O\Big(\min\Big(1,\frac{1}{H\|\frac{q_1x}{n_1}-\frac{r_2}{q_2}\|}\Big)\Big),$$

$$G_{21}'(x;H)=\sum_{\substack{n_2\le q_2\sqrt{T}\\ n_2\equiv r_2 \,(\mathrm{mod}\ q_2)}} O\Big(\min\Big(1,\frac{1}{H\|\frac{q_2x}{n_2}-\frac{r_1}{q_1}\|}\Big)\Big).$$

*Proof of Lemma 2.1.* From (1.4), we write

$$\Delta(q_1q_2x;r_1,q_1,r_2,q_2)=F_1+F_2+O(1), \tag{2.6}$$

where

$$F_1(x)=-\sum_{\substack{n_1\le q_1\sqrt{x}\\ n_1\equiv r_1 \,(\mathrm{mod}\, q_1)}}\psi\big(\frac{q_1x}{n_1}-\frac{r_2}{q_2}\big),\quad F_2(x)=-\sum_{\substack{n_2\le q_2\sqrt{x}\\ n_2\equiv r_2 \,(\mathrm{mod}\, q_2)}}\psi\big(\frac{q_2x}{n_2}-\frac{r_1}{q_1}\big).$$

The rest of the proof is almost the same as that for Lemma 3.1 in Liu [9].□

## 3. Preliminaries

We introduce some notations first. Set

$$E(n;H,J)=\{\{h,l\}\mid hl=n,1\le h\le H,h\le l\le 2^{J+1}h\}.$$

Denote

$$\begin{aligned}\Delta d_2(n;a_1,q_1,a_2,q_2)=&d(n;a_1,q_1,a_2,q_2)+d(n;-a_1,q_1,a_2,q_2)\\&-d(n;a_1,q_1,-a_2,q_2)-d(n;-a_1,q_1,a_2,q_2),\end{aligned}$$

$$\begin{aligned}&\Delta d_{2,1}(n,H,J;a_1,q_1,a_2,q_2)\\&=\sum_{\substack{\{h,l\}\in E(n;H,J)\\ h\equiv a_2 \,(\mathrm{mod}\ q_2)\\ l\equiv a_1 \,(\mathrm{mod}\ q_1)}}{}'1+\sum_{\substack{\{h,l\}\in E(n;H,J)\\ h\equiv a_2 \,(\mathrm{mod}\ q_2)\\ l\equiv -a_1 \,(\mathrm{mod}\ q_1)}}{}'1-\sum_{\substack{\{h,l\}\in E(n;H,J)\\ h\equiv -a_2 \,(\mathrm{mod}\ q_2)\\ l\equiv a_1 \,(\mathrm{mod}\ q_1)}}{}'1-\sum_{\substack{\{h,l\}\in E(n;H,J)\\ h\equiv -a_2 \,(\mathrm{mod}\ q_2)\\ l\equiv -a_1 \,(\mathrm{mod}\ q_1)}}{}'1,\end{aligned}$$

$$\begin{aligned}&\Delta d_{2,2}(n,H,J;a_1,q_1,a_2,q_2)\\ &=\sum_{\substack{\{h,l\}\in E(n;H,J)\\ h\equiv a_1\,(\mathrm{mod}\ q_1)\\ l\equiv a_2\,(\mathrm{mod}\ q_2)}}{}'\ 1+\sum_{\substack{\{h,l\}\in E(n;H,J)\\ h\equiv -a_1\,(\mathrm{mod}\ q_1)\\ l\equiv a_2\,(\mathrm{mod}\ q_2)}}{}'\ 1-\sum_{\substack{\{h,l\}\in E(n;H,J)\\ h\equiv a_1\,(\mathrm{mod}\ q_1)\\ l\equiv -a_2\,(\mathrm{mod}\ q_2)}}{}'\ 1-\sum_{\substack{\{h,l\}\in E(n;H,J)\\ h\equiv -a_1\,(\mathrm{mod}\ q_1)\\ l\equiv -a_2\,(\mathrm{mod}\ q_2)}}{}'\ 1.\end{aligned}$$

Set $\mathbf{1}=\{0,1\}$, $\boldsymbol{i}=(i_1,\cdots,i_{k-1})\in\mathbf{1}^{k-1}$, $\boldsymbol{n}=(n_1,\cdots,n_k)\in\mathbb{N}^k$. Define

$$\begin{aligned}&|\boldsymbol{i}|=i_1+\cdots+i_{k-1},\\ &\alpha(\boldsymbol{n};\boldsymbol{i})=\sqrt{n_1}+(-1)^{i_1}\sqrt{n_2}+\cdots+(-1)^{i_{k-1}}\sqrt{n_k},\\ &\beta(\boldsymbol{i})=\frac{3\pi}{4}+(-1)^{i_1}\frac{3\pi}{4}+\cdots+(-1)^{i_{k-1}}\frac{3\pi}{4}=(k-2|\boldsymbol{i}|)\frac{3\pi}{4},\\ &\mathcal{S}(k,\upsilon)=\{\{n_1,\dots n_k\}\mid \sqrt{n_1}+\cdots+\sqrt{n_\upsilon}=\sqrt{n_{\upsilon+1}}+\cdots+\sqrt{n_k}\}.\end{aligned}$$

Suppose $A_0>2$. Denote

$$K_0:=\min\{n\in\mathbb{N}: n\geq A_0, 2|n\},\qquad\qquad s(k):=2^{k-2}+\frac{k-6}{4}.$$

To prove the theorems, we need the following definition and Lemmas.

**Definition 3.1.** *( See eg.[7, p.73] ) Let $A>\frac{1}{2}$, $B\geq 1$, and $f(x)\in C^5[B,2B]$ be real valued function. Assume the derivatives of $f(x)$ for $B\leq x\leq 2B$ satisfy*

$$AB^{1-r}\ll|f^{(r)}(x)|\ll AB^{1-r},\quad r=1,2,3,4,$$

*where the $\ll$ constants depend on $r$ alone. Consider*

$$S_f(d)=\sum_{B<n\leq B+d}e(f(n)),\quad 1<d\leq B.$$

*The pair of nonnegative real numbers $(\kappa,\lambda)$ will be called an exponent pair, if $0\leq\kappa\leq\frac{1}{2}\leq\lambda\leq 1$ and*

$$S_f(d)\ll A^\kappa B^\lambda$$

*holds uniformly for $f$ and $d$.*

**Lemma 3.1.** *( See [15, p. 210] ) Let $H\geq 2$ be any real number. Then*

$$\psi(u)=\sum_{1\leq|h|\leq H}a(h)e(hu)+O\big(\sum_{1\leq|h|\leq H}b(h)e(hu)\big),$$

*where $a(h)$ and $b(h)$ are functions such that $a(h)\ll\frac{1}{|h|}$ and $b(h)\ll\frac{1}{H}$.*

**Lemma 3.2.** *( See [12, Theorem 2.2.2)] ) Let $A_1,\dots,A_5$ be absolute positive constants, $f(x)\in C^3[a,b]$ and $g(x)\in C^1[a,b]$ be $\mathbb{C}$ valued functions satisfying*

$$\begin{aligned}&\frac{A_1}{R}\leq|f''(x)|\leq\frac{A_1}{R},&&|f'''(x)|\leq\frac{A_3}{RU},\ U\geq 1,\\ &|g(x)|\leq A_4G,&&|g'(x)|\leq\frac{A_5G}{U_1},\ \ U_1\geq 1.\end{aligned}$$

*Suppose $\alpha \le f'(x) \le \beta$ for $a \le x \le b$. Then*

$$\sum_{a<x\le b} g(n)e\big(f(n)\big) = \sum_{\alpha\le u\le\beta} b_u \frac{g(n_u)}{\sqrt{f''(n_u)}} e\big(f(n_u) - un_u + \frac{1}{8}\big) + O\big(G\log(\beta-\alpha+2) + G(b-a+R)(U^{-1}+U_1^{-1})\big) + O\Big(G\min\Big\{\sqrt{R}, \max\Big(\frac{1}{\langle\alpha\rangle}, \frac{1}{\langle\beta\rangle}\Big)\Big\}\Big),$$

*where $n_u$ is the solution of $f'(n) = u$,*

$$\langle t\rangle = \begin{cases} \|t\|, & \text{if } t \text{ is not an integer,} \\ \beta-\alpha, & \text{if } t \text{ is an integer,} \end{cases}$$

$$b_u = \begin{cases} 1, & \text{if } \alpha < u < \beta, \text{ or } \alpha, \ \beta \text{ are not integers,} \\ \frac{1}{2}, & \text{if } u = \alpha \text{ or } \beta \text{ is an integer,} \end{cases}$$

$$\sqrt{f''} = \begin{cases} \sqrt{f''}, & \text{if } f'' > 0, \\ i\sqrt{f''}, & \text{if } f'' < 0. \end{cases}$$

**Lemma 3.3.** (*Halász-Montgomery inequality*) (*see e.g., Ivić[7]*) *Let $\xi$, $\varphi_1,\dots,$ $\varphi_R$ are arbitrary vectors in an inner-product vector space over $\mathbb{C}$, where $(a,b)$ will be the notion for the inner product and $\|a\|^2 = (a,a)$. Then*

$$\sum_{r\le R} \big|(\xi,\varphi_r)\big|^2 \le \|\xi\|^2 \max_{r\le R} \sum_{s\le R} \big|(\varphi_r,\varphi_s)\big|.$$

**Lemma 3.4.** (*See [17, eq.(4.6)]* ) *Suppose $k \ge 2$ is an integer, $y$ is a large positive number, $\boldsymbol{i} = (i_1,\cdots,i_{k-1}) \in \mathbf{1}^{k-1}$, and $\boldsymbol{n} = (n_1,\cdots,n_k) \in \mathbb{N}^k$. Then*

$$\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \sum_{\substack{n_i\le y \\ 1\le i\le k \\ \alpha(\boldsymbol{n};\boldsymbol{i})\ne 0}} \frac{d(n_1)\cdots d(n_k)}{(n_1\cdots n_k)^{\frac{3}{4}}\alpha(\boldsymbol{n};\boldsymbol{i})} \ll y^{s(k)+\varepsilon}.$$

**Lemma 3.5.** ( *See [17, Lemma 3.1]* ) *Suppose $f(n) : \mathbb{N} \to \mathbb{R}$ is a function with $f(n) \ll n^{\varepsilon'}$, $k \ge 2$ is an integer, $1 \le \upsilon < k$ is a fixed integer, $y > 1$ is large. Define*

$$s_{k,\upsilon}(f;y) := \sum_{\substack{\{n_1,n_2,\cdots,n_k\}\in\mathcal{S}(k,\upsilon) \\ n_1,n_2,\cdots,n_k\le y}} \frac{f(n_1)f(n_2)\cdots f(n_k)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}, \quad 1\le\upsilon<k,$$

$$s_{k,\upsilon}(f) := \sum_{\{n_1,n_2,\cdots,n_k\}\in\mathcal{S}(k,\upsilon)} \frac{f(n_1)f(n_2)\cdots f(n_k)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}, \quad 1\le\upsilon<k.$$

*Then for any $\varepsilon > 0$, $1 \le \upsilon < k$, we have*

$$s_{k,\upsilon}(f) \ll 1,$$
$$|s_{k,\upsilon}(f) - s_{k,\upsilon}(f;y)| \ll y^{-\frac{1}{2}+\varepsilon}.$$

**Lemma 3.6.** ( *Hilbert's inequality* )( *See e.g.[13]* ) *Let $x_1 < x_2 < \cdots < x_n$ be a sequence of real numbers. If there exists $\delta > 0$, such that $\min\limits_{s\neq r} |x_r - x_s| \geq \delta_r \geq \delta > 0 (1 \leq r \leq n)$, then there exists an absolute constant $C$, such that*

$$\left| \sum_{s\neq r} u_r \bar{u_s} (x_r - x_s)^{-1} \right| \leq C \sum_r {\delta_r}^{-1} |u_r|^2 ,$$

*for arbitrary complex numbers $u_1, u_2, \cdots, u_n$.*

## 4. Proofs of Theorem 1.1

We follow the approach of Zhai [18], and first present a large value estimate of $\Delta(q_1q_2x; r_1, q_1, r_2, q_2)$.

*4.1. Large value estimate of $\Delta(q_1q_2x; r_1, q_1, r_2, q_2)$*

We shall prove the following

**Theorem 4.1.** *Suppose $T \gg (q_1q_2)^{\varepsilon}$. Let $\frac{T}{2} \leq x_1 < \cdots < x_M \leq T$ satisfy $|\Delta(q_1q_2x_s; r_1, q_1, r_2, q_2)| \gg V$ $(s = 1, \ldots, M)$ and $|x_i - x_j| \geq V \gg T^{\frac{7}{32}} \mathcal{L}^4 (i \neq j)$. Then*

$$M \ll TV^{-3}\mathcal{L}^9 + T^{\frac{15}{4}} V^{-12} \mathcal{L}^{41}.$$

*Proof of Theorem 4.1.* Let $T_0 > V$ be a parameter to be determined and $I$ be any subinterval of $[\frac{T}{2}, T]$ of length not exceeding $T_0$. Without loss of generality, we may assume $G = I \cap \{x_1, \ldots, x_M\} = \{x_1, \ldots, x_{M_0}\}$.

From (2.6), let $J_0 = \left[\frac{\mathcal{L} + 2\log\mathcal{L} - 2\log V}{\log 2}\right]$ and

$$\Psi_1(x; j) = \sum_{2^{-\frac{j+1}{2}} q_1\sqrt{x} \leq k_1q_1 + r_1 \leq 2^{-\frac{j}{2}} q_1\sqrt{x}} \psi\Big(\frac{q_1x}{k_1q_1 + r_1} - \frac{r_2}{q_2}\Big),$$

$$\Psi_2(x; j) = \sum_{2^{-\frac{j+1}{2}} q_2\sqrt{x} \leq k_2q_2 + r_2 \leq 2^{-\frac{j}{2}} q_2\sqrt{x}} \psi\Big(\frac{q_2x}{k_2q_2 + r_2} - \frac{r_1}{q_1}\Big).$$

Then

$$\Delta(q_1q_2x; r_1, q_1, r_2, q_2) = -\sum_{j=0}^{J_0} \Psi_1(x; j) - \sum_{j=0}^{J_0} \Psi_2(x; j) + O\Big(\frac{V}{\mathcal{L}}\Big). \tag{4.1}$$

By Cauchy's inequality we get

$$\Delta^2(q_1q_2x; r_1, q_1, r_2, q_2) \ll \mathcal{L} \sum_{j=0}^{J_0} \Psi_1^2(x; j) + \mathcal{L} \sum_{j=0}^{J_0} \Psi_2^2(x; j) + O\Big(\frac{V^2}{\mathcal{L}^2}\Big),$$

which implies that for $x \in G$

$$\Delta^2(q_1q_2x; r_1, q_1, r_2, q_2) \ll \mathcal{L}\sum_{j=0}^{J_0}\Psi_1^2(x; j) + \mathcal{L}\sum_{j=0}^{J_0}\Psi_2^2(x; j).$$

Denote

$$G_1 := \Big\{x_s \in G \;\Big|\; \sum_{j=0}^{J_0}\Psi_1^2(x_s; j) \geq \sum_{j=0}^{J_0}\Psi_2^2(x_s; j)\Big\}, \qquad M_1 =: \#G_1;$$

$$G_2 := \Big\{x_s \in G \;\Big|\; \sum_{j=0}^{J_0}\Psi_2^2(x_s; j) \geq \sum_{j=0}^{J_0}\Psi_1^2(x_s; j)\Big\}, \qquad M_2 =: \#G_2.$$

We first estimate $M_1$. Assume that $G_1 = \{x_1, \ldots, x_{M_1}\}$. Then we have

$$\Delta^2(q_1q_2x_s; r_1, q_1, r_2, q_2) \ll \mathcal{L}\sum_{j=0}^{J_0}\Psi_1^2(x_s; j), \quad \text{for } 1 \leq s \leq M_1.$$

Summing over the set $G_1$ we get

$$\begin{aligned} M_1V^2 &\ll \sum_{s\leq M_1}\Delta^2(q_1q_2x_s; r_1, q_1, r_2, q_2) \qquad (4.2)\\ &\ll\mathcal{L}\sum_{j=0}^{J_0}\sum_{s\leq M_1}\Psi_1^2(x_s; j) \ll \mathcal{L}^2\sum_{s\leq M_1}\Psi_1^2(x_s; j), \end{aligned}$$

for some fixed $j$ with $0 \leq j \leq J_0$. Let $2 \leq H < T$ be a parameter to be determined later. From Lemma 3.1 and Lemma 3.2, we have

$$\Psi_1(x_s; j) \ll \Big|\sum_{1\leq h\leq H} c(h) \sum_{\substack{k_1\\ 2^{-\frac{j+1}{2}}q_1\sqrt{x_s}\leq k_1q_1+r_1\leq 2^{-\frac{j}{2}}q_1\sqrt{x_s}}} e\Big(-h\big(\frac{q_1x_s}{k_1q_1+r_1} - \frac{r_2}{q_2}\big)\Big)\Big|,$$

where $c(h)$ is some function such that $c(h) \ll \frac{1}{h}$. By using Lemma 3.2, we get

$$\Psi_1(x_s; j) \ll \frac{x^{\frac{1}{4}}}{\sqrt{2}}\sum_{1\leq h\leq H} c(h)\sum_{2^jh\leq l\leq 2^{j+1}h}\frac{b_lh^{\frac{1}{4}}}{l^{\frac{3}{4}}}e\big(-2\sqrt{hlx_s}+\frac{hr_2}{q_2}+\frac{lr_1}{q_1}\big)+O(\mathcal{L}^2). \qquad (4.3)$$

Inserting (4.3) into (4.2) yields

$$\begin{aligned} M_1V^2 &\ll \mathcal{L}^2T^{\frac{1}{2}}\sum_{s\leq M_1}\Big|\sum_{1\leq h\leq H} c(h)\sum_{2^jh\leq l\leq 2^{j+1}h}\frac{b_lh^{\frac{1}{4}}}{l^{\frac{3}{4}}}e\big(-2\sqrt{hlx_s}+\frac{hr_2}{q_2}+\frac{lr_1}{q_1}\big)\Big|^2+M_1\mathcal{L}^6\\ &\ll \mathcal{L}^2T^{\frac{1}{2}}\sum_{s\leq M_1}\Big|\sum_{1\leq n\leq 2^{j+1}H^2}\sum_{\substack{n=hl\\ 1\leq h\leq H\\ 2^jh\leq l\leq 2^{j+1}h}}\frac{b_lhc(h)}{n^{\frac{3}{4}}}e\big(-2\sqrt{hlx_s}+\frac{hr_2}{q_2}+\frac{lr_1}{q_1}\big)\Big|^2. \end{aligned}$$

Let

$$\gamma(n;H,j)=\sum_{\substack{n=hl\\ 1\le h\le H\\ 2^jh\le l\le 2^{j+1}h}} b_l hc(h)e\big(\frac{hr_2}{q_2}+\frac{lr_1}{q_1}\big).$$

Then it is easy to see that $\gamma(n;H,j)\ll d(n)$, and

$$M_1V^2\ll\mathcal{L}^2T^{\frac12}\sum_{s\le M_1}\Big|\sum_{1\le n\le 2^{j+1}H^2}\frac{\gamma(n;H,j)}{n^{\frac34}}e(-2\sqrt{nx_s})\Big|^2.$$

By a splitting argument and Cauchy's inequality, we get

$$\begin{aligned}M_1V^2\ll&\mathcal{L}^2T^{\frac12}\sum_{s\le M_1}\Big|\sum_{\nu}\sum_{2^{j-v}H^2<n\le 2^{j+1-v}H^2}\frac{\gamma(n;H,j)}{n^{\frac34}}e(-2\sqrt{nx_s})\Big|^2\\ \ll&\mathcal{L}^3T^{\frac12}\sum_{\nu}\sum_{s\le M_1}\Big|\sum_{2^{j-v}H^2<n\le 2^{j+1-v}H^2}\frac{\gamma(n;H,j)}{n^{\frac34}}e(-2\sqrt{nx_s})\Big|^2\\ \ll&\mathcal{L}^4T^{\frac12}\sum_{s\le M_1}\Big|\sum_{2^{j-v}H^2<n\le 2^{j+1-v}H^2}\frac{\gamma(n;H,j)}{n^{\frac34}}e(-2\sqrt{nx_s})\Big|^2\end{aligned}$$

for some $0\le\nu\ll\mathcal{L}$. Let $N_1=2^{j-v}H^2$. Take $\xi=\{\xi_n\}_{n=1}^\infty$ with $\xi_n=\gamma(n;H,j)n^{-\frac34}$ for $N_1<n\le 2N_1$ and zero otherwise, and $\varphi_s=\{\varphi_{s,n}\}_{n=1}^\infty$ with $\varphi_{s,n}=e(-2\sqrt{nx_s})$ for $N_1<n\le 2N_1$ and zero otherwise. Then

$$\begin{aligned}\sum_{s\le M_1}(\xi,\varphi_s)^2=&\sum_{s\le M_1}\Big|\sum_{N_1<n\le 2N_1}\frac{\gamma(n;H,j)}{n^{\frac34}}e(-2\sqrt{nx_s})\Big|^2,\\ (\varphi_{s_1},\varphi_{s_2})=&\sum_{N_1<n\le 2N_1}e\big(2\sqrt{n}(\sqrt{x_{s_2}}-\sqrt{x_{s_1}})\big),\\ \|\xi\|^2=&\sum_{N_1<n\le 2N_1}\frac{|\gamma(n;H,j)|^2}{n^{\frac32}}\ll\sum_{N_1<n\le 2N_1}\frac{d^2(n)}{n^{\frac32}}\ll N_1^{-\frac12}\mathcal{L}^3,\end{aligned}$$

where we used the estimate $\sum_{n\le N}d^2(n)\ll N\log^3 N$. Thus by Lemma 3.3, we get

$$\begin{aligned}M_1V^2\ll&T^{\frac12}N_1^{-\frac12}\mathcal{L}^7\max_{s_1\le M_1}\sum_{s_2\le M_1}\Big|\sum_{N_1<n\le 2N_1}e\big(2\sqrt{n}(\sqrt{x_{s_2}}-\sqrt{x_{s_1}})\big)\Big| \qquad (4.4)\\ =&T^{\frac12}N_1^{\frac12}\mathcal{L}^7+T^{\frac12}N_1^{-\frac12}\mathcal{L}^7\max_{s_1\le M_1}\sum_{\substack{s_2\le M_1\\ s_2\ne s_1}}\Big|\sum_{N_1<n\le 2N_1}e\big(2\sqrt{n}(\sqrt{x_{s_2}}-\sqrt{x_{s_1}})\big)\Big|.\end{aligned}$$

By the Kuz'min-Landau inequality and the exponent pair $(\frac{4}{18},\frac{11}{18})$ (see eg. [7,

p.77]), we have

$$\begin{aligned}\sum_{N_1<n\le 2N_1} e\big(2\sqrt{n}(\sqrt{x_{s_2}}-\sqrt{x_{s_1}})\big) \ll&\frac{\sqrt{N_1}}{|\sqrt{x_{s_2}}-\sqrt{x_{s_1}}|}+\Big(\frac{|\sqrt{x_{s_2}}-\sqrt{x_{s_1}}|}{\sqrt{N_1}}\Big)^{\frac{4}{18}}N_1^{\frac{11}{18}}\\ \ll&\frac{\sqrt{N_1T}}{|x_{s_2}-x_{s_1}|}+\Big(\frac{|x_{s_2}-x_{s_1}|}{\sqrt{N_1T}}\Big)^{\frac{4}{18}}N_1^{\frac{11}{18}}\\ \ll&\frac{\sqrt{N_1T}}{|s_2-s_1|V}+T^{-\frac{1}{9}}T_0^{\frac{2}{9}}N_1^{\frac{1}{2}},\end{aligned}$$

in view of $|x_{s_2}-x_{s_1}|\le T_0$. Combining this with (4.4), we get

$$\begin{aligned}M_1V^2\ll&T^{\frac12}N_1^{\frac12}\mathcal{L}^7+TV^{-1}\mathcal{L}^7\max_{s_1\le M_1}\sum_{\substack{s_2\le M_1\\ s_2\ne s_1}}\frac{1}{|s_2-s_1|}+M_0T^{\frac{7}{18}}T_0^{\frac29}\mathcal{L}^7 \qquad (4.5)\\ \ll&T^{\frac12}N_1^{\frac12}\mathcal{L}^7+TV^{-1}\mathcal{L}^8+M_0T^{\frac{7}{18}}T_0^{\frac29}\mathcal{L}^7.\end{aligned}$$

Take $H=T^{\frac12}V^{-1}2^{-\frac{j}{2}}\mathcal{L}^2$, $T_0=T^{-\frac74}V^9\mathcal{L}^{-32}$. Then $N_1=TV^{-2}2^{-\nu}\mathcal{L}^4$, and $T_0\gg V$ if $V\gg T^{\frac{7}{32}}\mathcal{L}^4$. Thus from (4.5), we get

$$M_1\ll TV^{-3}\mathcal{L}^9.$$

For $M_2$, by symmetry, we can get the same estimate. Noting that $M_1+M_2\ge M_0$, we have

$$M_0\ll TV^{-3}\mathcal{L}^9.$$

Dividing the interval $[\frac{T}{2},T]$ into $O\big(1+\frac{T}{2T_0}\big)$ subintervals of length not exceeding $T_0=T^{-\frac74}V^9\mathcal{L}^{-32}$, we see

$$M\ll M_0\big(1+\frac{T}{2T_0}\big)\ll TV^{-3}\mathcal{L}^9+T^{\frac{15}{4}}V^{-12}\mathcal{L}^{41}.$$

This completes the proof of Theorem 4.1. □

*4.2. Proof of Theorem 1.1*

For $A=0$, this is trivial. For $0<A\le 2$, it follows from (2.4) and Hölder's inequality. So it suffices to prove it for $A>2$.

With a similar argument to (13.70) of Ivić, we see

$$\int_{\frac{T}{2}}^{T}|\Delta(q_1q_2x;r_1,q_1,r_2,q_2)|^Adx\ll T^{1+\frac{A}{4}}\mathcal{L}^{4A}+\sum_V V\sum_{s\le N_V}|\Delta(q_1q_2x_s;r_1,q_1,r_2,q_2)|^A,$$

where $T^{\frac14}\mathcal{L}^4\le V\le T^{\frac{131}{416}}\mathcal{L}^4$, $V<|\Delta(q_1q_2x_s;r_1,q_1,r_2,q_2)|\le 2V$, $s=1,\dots,N_V$, $|x_{s_1}-x_{s_2}|\ge V$ for $s_1\ne s_2\le N_V$, and the number of $V$ is $\ll\mathcal{L}$. By Theorem 4.1, we get

$$V\sum_{s\le N_V}|\Delta(q_1q_2x_s;r_1,q_1,r_2,q_2)|^A\ll N_VV^{A+1}\ll TV^{A-2}\mathcal{L}^9+T^{\frac{15}{4}}V^{A-11}\mathcal{L}^{41}.$$

Thus we have

$$\int_{\frac{T}{2}}^{T} |\Delta(q_1q_2x;r_1,q_1,r_2,q_2)|^A dx \ll T^{1+\frac{A}{4}}\mathcal{L}^{4A} + TV^{A-2}\mathcal{L}^{10} + T^{\frac{15}{4}}V^{A-11}\mathcal{L}^{42},$$

for some fixed $T^{\frac{1}{4}}\mathcal{L}^4 \le V \le T^{\frac{131}{416}}\mathcal{L}^4$, which implies

$$\begin{aligned}
&\int_{\frac{T}{2}}^{T} |\Delta(q_1q_2x;r_1,q_1,r_2,q_2)|^A dx \\
&\ll \begin{cases} T^{1+\frac{A}{4}}\mathcal{L}^{4A} + T^{1+\frac{131}{416}(A-2)}\mathcal{L}^{4A+2} + T^{\frac{15}{4}+\frac{1}{4}(A-11)}\mathcal{L}^{4A-2}, & \text{if } 2 < A \le 11, \\ T^{1+\frac{A}{4}}\mathcal{L}^{4A} + T^{1+\frac{131}{416}(A-2)}\mathcal{L}^{4A+2} + T^{\frac{15}{4}+\frac{131}{416}(A-11)}\mathcal{L}^{4A-2}, & \text{if } A > 11, \end{cases} \\
&\ll \begin{cases} T^{1+\frac{A}{4}}\mathcal{L}^{4A}, & \text{if } 2 < A \le \frac{262}{27}, \\ T^{\frac{131}{416}A+\frac{154}{416}}\mathcal{L}^{4A+2}, & \text{if } A > 11. \end{cases}
\end{aligned}$$

Thus we obtain the estimates for integrals over intervals of the form $[2^{-j}T, 2^{1-j}T]$, $1 \le j \ll \log T$, by summing over which, we get Theorem 1.1. □

## 5. Proofs of Theorem 1.2 and Theorem 1.3

Suppose $x \ge 1, 1 \le a_i \le q_i$ and $(a_i, q_i) = 1$ for $i = 1, 2$. Since periods of both $\cos(2\pi m\theta_1)$ and $\sin(2\pi n\theta_2)$ are 1, we have

$$\begin{aligned}
S\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big) &= \sideset{}{'}\sum_{mn\le q_1q_2x} \cos\Big(2\pi m\frac{a_1}{q_1}\Big)\sin\Big(2\pi n\frac{a_2}{q_2}\Big) \\
&= \sum_{r_1=1}^{q_1}\sum_{r_2=1}^{q_2} \sideset{}{'}\sum_{\substack{mn\le q_1q_2x \\ m\equiv r_1 \,(\mathrm{mod}\ q_1) \\ n\equiv r_2 \,(\mathrm{mod}\ q_2)}} \cos\Big(2\pi r_1\frac{a_1}{q_1}\Big)\sin\Big(2\pi r_2\frac{a_2}{q_2}\Big) \\
&= \sum_{r_1=1}^{q_1}\sum_{r_2=1}^{q_2} \cos\Big(2\pi r_1\frac{a_1}{q_1}\Big)\sin\Big(2\pi r_2\frac{a_2}{q_2}\Big) \sideset{}{'}\sum_{\substack{mn\le q_1q_2x \\ m\equiv r_1 \,(\mathrm{mod}\ q_1) \\ n\equiv r_2 \,(\mathrm{mod}\ q_2)}} 1.
\end{aligned}$$

Substituting (1.4) into this formula, we obtain

$$\begin{aligned}
S\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big) =& \big(x\log x - x\big)\sum_{r_1=1}^{q_1}\cos\Big(2\pi r_1\frac{a_1}{q_1}\Big)\sum_{r_2=1}^{q_2}\sin\Big(2\pi r_2\frac{a_2}{q_2}\Big) \\
&- x\sum_{r_1=1}^{q_1}\cos\Big(2\pi r_1\frac{a_1}{q_1}\Big)\frac{\Gamma'}{\Gamma}\Big(\frac{r_1}{q_1}\Big)\sum_{r_2=1}^{q_2}\sin\Big(2\pi r_2\frac{a_2}{q_2}\Big) \\
&- x\sum_{r_2=1}^{q_2}\sin\Big(2\pi r_2\frac{a_2}{q_2}\Big)\frac{\Gamma'}{\Gamma}\Big(\frac{r_2}{q_2}\Big)\sum_{r_1=1}^{q_1}\cos\Big(2\pi r_1\frac{a_1}{q_1}\Big) \\
&+ \sum_{r_1=1}^{q_1}\sum_{r_2=1}^{q_2}\cos\big(2\pi r_1\frac{a_1}{q_1}\big)\sin\big(2\pi r_2\frac{a_2}{q_2}\big)\Delta(q_1q_2x;r_1,q_1,r_2,q_2).
\end{aligned} \tag{5.1}$$

For $(a,q)=1$ and $\theta$ an arbitrary constant, it is easy to see that

$$\sum_{r=1}^{q}\sin(2\pi r\frac{a}{q}+\theta)=\sum_{r=1}^{q}\cos(2\pi r\frac{a}{q}+\theta)=0, \tag{5.2}$$

which implies

$$S\big(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2}\big)=\sum_{r_1=1}^{q_1}\sum_{r_2=1}^{q_2}\cos\big(2\pi\frac{r_1a_1}{q_1}\big)\sin\big(2\pi\frac{r_2a_2}{q_2}\big)\Delta(q_1q_2x;r_1,q_1,r_2,q_2). \tag{5.3}$$

Using the estimate of $\Delta(x;r_1,q_1,r_2,q_2)$ shown by (2.1), and Müller and Nowak's (2.3), we complete the proof of Theorem 1.2 and Theorem 1.3.

## 6. Proofs of Theorem 1.4

### *6.1. An expression of* $S^k(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2})$

From (5.3), (2.5), noting the fact that $T\gg(q_1q_2)^{\varepsilon'}$, we have

$$\begin{aligned}&S\big(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2}\big)\\ =&R_0(x;y)+R_{12}(x;y,H)+R_{21}(x;y,H)+G_{12}(x;H)+G_{21}(x;H)+O\big(q_1q_2\mathcal{L}^3\big).\end{aligned}$$

Here

$$R_0(x;y)=\frac{x^{\frac14}}{\sqrt2\pi}\sum_{r_1=1}^{q_1}\sum_{r_2=1}^{q_2}\cos\big(2\pi r_1\frac{a_1}{q_1}\big)\sin\big(2\pi r_2\frac{a_2}{q_2}\big)\sum_{n\le y}\frac{\tau(n,x)}{n^{\frac34}}, \tag{6.1}$$

$$R_{12}(x;y,H)=\frac{x^{\frac14}}{\sqrt2\pi}\sum_{r_1=1}^{q_1}\sum_{r_2=1}^{q_2}\cos\big(2\pi r_1\frac{a_1}{q_1}\big)\sin\big(2\pi r_2\frac{a_2}{q_2}\big)\sum_{y<n\le2^{J+1}H^2}\frac{\tau_{12}(n,x;H)}{n^{\frac34}},$$

$$R_{21}(x;y,H)=\frac{x^{\frac14}}{\sqrt2\pi}\sum_{r_1=1}^{q_1}\sum_{r_2=1}^{q_2}\cos\big(2\pi r_1\frac{a_1}{q_1}\big)\sin\big(2\pi r_2\frac{a_2}{q_2}\big)\sum_{y<n\le2^{J+1}H^2}\frac{\tau_{21}(n,x;H)}{n^{\frac34}},$$

$$G_{12}(x;H)=O\Big(q_2\sum_{n_1\le q_1\sqrt T}\min\Big(1,\frac{1}{H\|\frac{q_1x}{n_1}-\frac{r_2}{q_2}\|}\Big)\Big), \tag{6.2}$$

$$G_{21}(x;H)=O\Big(q_1\sum_{n_2\le q_2\sqrt T}\min\Big(1,\frac{1}{H\|\frac{q_2x}{n_2}-\frac{r_1}{q_1}\|}\Big)\Big).$$

Denote $R_0=R_0(x;y)$, $R_1=R_{12}(x;y,H)$, $R_2=R_{21}(x;y,H)$, $G_{12}=G_{12}(x;H)$, $G_{21}=G_{21}(x;H)$, $R=R_0+R_1+R_2$, and $G=G_{12}+G_{21}$. Then

$$S\big(q_1q_2x;\frac{a_1}{q_1},\frac{a_2}{q_2}\big)=R+G+O\big(q_1q_2\mathcal{L}^3\big).$$

Let $k \geq 2$ be a fixed integer. By the elementary formula $(a+b)^k = a^k + O(|a|^{k-1}|b| + |b|^k)$, we get

$$S^k\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big) = R^k + O\big(|R|^{k-1}G + |R|^{k-1}q_1q_2\mathcal{L}^3\big) + O\big(G^k + (q_1q_2\mathcal{L}^3)^k\big). \tag{6.3}$$

The expression of $R_0$ given by (6.1) shows that

$$\begin{aligned} R_0 =& \frac{x^{\frac{1}{4}}}{\sqrt{2}\pi} \sum_{n\leq y} n^{-\frac{3}{4}} \sum_{n=hl} \sum_{r_1=1}^{q_1} \sum_{r_2=1}^{q_2} \cos\big(2\pi r_1 \frac{a_1}{q_1}\big) \sin\big(2\pi r_2 \frac{a_2}{q_2}\big) \\ & \times \cos\Big(4\pi\sqrt{nx} - 2\pi\Big(\frac{hr_2}{q_2} + \frac{lr_1}{q_1} + \frac{1}{8}\Big)\Big). \end{aligned} \tag{6.4}$$

Set

$$S_0 = \sum_{r_1=1}^{q_1} \cos\big(2\pi r_1 \frac{a_1}{q_1}\big) \cos\Big(4\pi\sqrt{nx} - 2\pi\Big(\frac{hr_2}{q_2} + \frac{lr_1}{q_1} + \frac{1}{8}\Big)\Big).$$

Then, clearly

$$\begin{aligned} S_0 = \frac{1}{2}\sum_{r_1=1}^{q_1} \Bigg\{ & \cos\Big(4\pi\sqrt{nx} - 2\pi\Big(\frac{hr_2}{q_2} + \frac{r_1}{q_1}(l+a_1) + \frac{1}{8}\Big)\Big) \\ & + \cos\Big(4\pi\sqrt{nx} - 2\pi\Big(\frac{hr_2}{q_2} + \frac{r_1}{q_1}(l-a_1) + \frac{1}{8}\Big)\Big)\Bigg\}. \end{aligned}$$

If $l \not\equiv \pm a_1 (\mathrm{mod}\ q_1)$, from (5.2), $S_0 = 0$.

If $q_1 > 2$, then for $l \equiv a_1 (\mathrm{mod}\ q_1)$ or $l \equiv -a_1 (\mathrm{mod}\ q_1)$, obviously they cannot be simultaneously true,

$$S_0 = \frac{1}{2}\sum_{r_1=1}^{q_1} \cos\Big(4\pi\sqrt{nx} - 2\pi\big(\frac{hr_2}{q_2} + \frac{1}{8}\big)\Big) = \frac{q_1}{2}\cos\Big(4\pi\sqrt{nx} - 2\pi\big(\frac{hr_2}{q_2} + \frac{1}{8}\big)\Big).$$

If $q_1 = 2$, then $a_1 = 1$, and $a_1 \equiv -a_1 (\mathrm{mod}\ q_1)$. Thus, for $l \equiv a_1 (\mathrm{or} -a_1)$ $(\mathrm{mod}\ q_1)$,

$$S_0 = q_1 \cos\Big(4\pi\sqrt{nx} - 2\pi\big(\frac{hr_2}{q_2} + \frac{1}{8}\big)\Big).$$

Summing up, we see

$$\begin{aligned} & \sum_{r_1=1}^{q_1} \cos\big(2\pi r_1 \frac{a_1}{q_1}\big) \cos\Big(4\pi\sqrt{nx} - 2\pi\big(\frac{hr_2}{q_2} + \frac{lr_1}{q_1} + \frac{1}{8}\big)\Big) \\ = & \begin{cases} \frac{q_1}{2}\cos\Big(4\pi\sqrt{nx} - 2\pi\big(\frac{hr_2}{q_2} + \frac{1}{8}\big)\Big), & q_1 > 2, l \equiv a_1, -a_1 \,(\mathrm{mod}\ q_1); \\ q_1 \cos\Big(4\pi\sqrt{nx} - 2\pi\big(\frac{hr_2}{q_2} + \frac{1}{8}\big)\Big), & q_1 = 2, l \equiv a_1 \,(\mathrm{mod}\ q_1); \\ 0, & \text{else.} \end{cases} \end{aligned} \tag{6.5}$$

By a similar argument, we get

$$\begin{aligned}&\sum_{r_2=1}^{q_2}\sin\big(2\pi r_2\frac{a_2}{q_2}\big)\cos\Big(4\pi\sqrt{nx}-2\pi\big(\frac{hr_2}{q_2}+\frac{1}{8}\big)\Big) \\ =&\begin{cases}\frac{q_2}{2}\cos\big(4\pi\sqrt{nx}-\frac{3\pi}{4}\big), & h\equiv a_2\,(\mathrm{mod}\ q_2);\\ -\frac{q_2}{2}\cos\big(4\pi\sqrt{nx}-\frac{3\pi}{4}\big), & h\equiv -a_2\,(\mathrm{mod}\ q_2);\\ 0, & \text{else.}\end{cases}\end{aligned} \tag{6.6}$$

Combining (6.4)-(6.6), we obtain

$$\begin{aligned}R_0=&\frac{x^{\frac{1}{4}}}{\sqrt{2}\pi}\times\frac{1}{2}\times\frac{1}{2}q_1q_2\sum_{n\le y}\frac{\cos\big(4\pi\sqrt{nx}-\frac{3\pi}{4}\big)}{n^{\frac{3}{4}}} \\ &\times\Big(\sum_{\substack{n=hl\\ h\equiv a_2\,(\mathrm{mod}\ q_2)\\ l\equiv a_1\,(\mathrm{mod}\ q_1)}}1+\sum_{\substack{n=hl\\ h\equiv a_2\,(\mathrm{mod}\ q_2)\\ l\equiv -a_1\,(\mathrm{mod}\ q_1)}}1-\sum_{\substack{n=hl\\ h\equiv -a_2\,(\mathrm{mod}\ q_2)\\ l\equiv a_1\,(\mathrm{mod}\ q_1)}}1-\sum_{\substack{n=hl\\ h\equiv -a_2\,(\mathrm{mod}\ q_2)\\ l\equiv -a_1\,(\mathrm{mod}\ q_1)}}1\Big)\\ =&\frac{q_1q_2x^{\frac{1}{4}}}{4\sqrt{2}\pi}\sum_{n\le y}\frac{\cos\big(4\pi\sqrt{nx}-\frac{3\pi}{4}\big)}{n^{\frac{3}{4}}}\Delta d_2(n;a_1,q_1,a_2,q_2).\end{aligned} \tag{6.7}$$

Similarly, we have

$$\begin{aligned}R_1=&\frac{q_1q_2x^{\frac{1}{4}}}{4\sqrt{2}\pi}\sum_{y<n\le 2^{J+1}H^2}\frac{\cos\big(4\pi\sqrt{nx}-\frac{3\pi}{4}\big)}{n^{\frac{3}{4}}}\Delta d_{2,1}(n,H,J;a_1,q_1,a_2,q_2),\\ R_2=&\frac{q_1q_2x^{\frac{1}{4}}}{4\sqrt{2}\pi}\sum_{y<n\le 2^{J+1}H^2}\frac{\cos\big(4\pi\sqrt{nx}-\frac{3\pi}{4}\big)}{n^{\frac{3}{4}}}\Delta d_{2,2}(n,H,J;a_1,q_1,a_2,q_2).\end{aligned} \tag{6.8}$$

*6.2. Upper bound of* $\int_{\frac{T}{2}}^{T}G^k dx\ \ (k\ge 1)$

From the expression of $G_{12}$ provided by (6.2), we have

$$\int_{\frac{T}{2}}^{T}G_{12}(x;H)dx\ll q_2\sum_{n_1\le q_1\sqrt{T}}\int_{\frac{T}{2}}^{T}\min\Big(1,\frac{1}{H\|\frac{q_1x}{n_1}-\frac{r_2}{q_2}\|}\Big)dx.$$

Let $u = \dfrac{q_1 x}{n_1} - \dfrac{l_2}{q_2}$. Then

$$\begin{aligned}\int_{\frac{T}{2}}^{T} G_{12}(x;H)dx \ll & q_2 \sum_{n_1 \le q_1\sqrt{T}} \int_{\frac{q_1 T}{2n_1}-\frac{r_2}{q_2}}^{\frac{q_1 T}{n_1}-\frac{r_2}{q_2}} \min\Big(1, \frac{1}{H\|u\|}\Big)\frac{n_1}{q_1}du \\ \ll & T q_2 \sum_{n_1 \le q_1\sqrt{T}} \int_0^1 \min\Big(1, \frac{1}{H\|u\|}\Big)du \\ \ll & q_1 q_2 T^{\frac{3}{2}} \int_0^{\frac{1}{2}} \min\Big(1, \frac{1}{Hu}\Big)du.\end{aligned} \tag{6.9}$$

Since

$$\int_0^{\frac{1}{2}} \min\Big(1, \frac{1}{Hu}\Big)du = \int_{\frac{1}{H}}^{\frac{1}{2}} \frac{1}{Hu}du + \int_0^{\frac{1}{H}} 1du \ll \frac{1}{H}\mathcal{L},$$

we obtain

$$\int_{\frac{T}{2}}^{T} G_{12}(x;H)dx \ll q_1 q_2 T^{\frac{3}{2}} H^{-1}\mathcal{L}.$$

For $\int_{\frac{T}{2}}^{T} G_{21}(x;H)dx$, this estimate still holds, which implies

$$\int_{\frac{T}{2}}^{T} G(x;H)dx \ll q_1 q_2 T^{\frac{3}{2}} H^{-1}\mathcal{L}.$$

Then by $G \ll q_1 q_2 T^{\frac{1}{2}}$, we get for $k \ge 1$,

$$\int_{\frac{T}{2}}^{T} G^k(x)dx \ll \big(q_1 q_2 T^{\frac{1}{2}}\big)^{k-1} \int_{\frac{T}{2}}^{T} G(x)dx \ll (q_1 q_2)^k T^{1+\frac{k}{2}} H^{-1}\mathcal{L}.$$

Taking $H = T^{\frac{A_0}{2}}$, we have the following

**Lemma 6.1.** *If $k \ge 1, k \in \mathbb{R}$, then*

$$\int_{\frac{T}{2}}^{T} G^k(x)dx \ll (q_1 q_2)^k T\mathcal{L}.$$

*6.3. Evaluation of the integral $\int_{\frac{T}{2}}^{T} R_0^2(x;y)dx$*

Now we go to evaluate the integral $\int_{\frac{T}{2}}^{T} R_0^2(x;y)dx$, which provides the main term in the asymptotic formula of $\int_{\frac{T}{2}}^{T} S^2(x, \frac{a_1}{q_1}, \frac{a_2}{q_2})dx$.

By the expression of $R_0$ given by (6.7), using the elementary formula

$$\cos a_1 \cos a_2 = \frac{1}{2}\cos(a_1 - a_2) + \frac{1}{2}\cos(a_1 + a_2),$$

we get

$$R_0^2(x;y)=\frac{(q_1q_2)^2}{64\pi^2}\big(S_1(x)+S_2(x)\big), \tag{6.10}$$

where

$$\begin{aligned}S_1(x)=&x^{\frac12}\sum_{n_1,n_2\le y}\frac{\Delta d_2(n_1;a_1,q_1,a_2,q_2)\Delta d_2(n_2;a_1,q_1,a_2,q_2)}{(n_1n_2)^{\frac34}}\\&\times\cos\big(4\pi\sqrt{x}(\sqrt{n_1}-\sqrt{n_2})\big),\\S_2(x)=&x^{\frac12}\sum_{n_1,n_2\le y}\frac{\Delta d_2(n_1;a_1,q_1,a_2,q_2)\Delta d_2(n_2;a_1,q_1,a_2,q_2)}{(n_1n_2)^{\frac34}}\\&\times\cos\big(4\pi\sqrt{x}(\sqrt{n_1}+\sqrt{n_2})-\frac{3\pi}{2}\big).\end{aligned}$$

Noticing that for $a,b>1$,

$$\int_a^b\cos(A\sqrt{t}+B)dt\ll\frac{\sqrt{b}+\sqrt{a}}{|A|}, \tag{6.11}$$

we have

$$\int_{\frac T2}^T x^{\frac k4}\cos(A\sqrt{x}+B)dx=\int_{\frac T2}^T x^{\frac k4}d\Big(\int_{\frac T2}^x\cos(A\sqrt{t}+B)dt\Big)\ll T^{\frac k4+\frac12}|A|^{-1}. \tag{6.12}$$

Hence, we get

$$\begin{aligned}\int_{\frac T2}^T S_2(x)dx=&\sum_{n_1,n_2\le y}\frac{\Delta d_2(n_1;a_1,q_1,a_2,q_2)\Delta d_2(n_2;a_1,q_1,a_2,q_2)}{(n_1n_2)^{\frac34}}\\&\times\int_{\frac T2}^T x^{\frac12}\cos\big(4\pi\sqrt{x}(\sqrt{n_1}+\sqrt{n_2})-\frac{3\pi}{2}\big)dx\\\ll&T\sum_{n_1,n_2\le y}\frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac34}(\sqrt{n_1}+\sqrt{n_2})}\\\ll&T\sum_{n_1,n_2\le y}\frac{d(n_1)d(n_2)}{n_1n_2}\ll T\log^4y,\end{aligned} \tag{6.13}$$

by using $\sqrt{n_1}+\sqrt{n_2}\ge2(n_1n_2)^{\frac14}$, and $\sum_{n\le N}\frac{d(n)}{n}\ll\log^2N$ in the last step.

Now we consider $\int_{\frac T2}^T S_1(x)dx$. Denote

$$S_1(x)=S_{11}(x)+S_{12}(x), \tag{6.14}$$

where

$$S_{11}(x) = x^{\frac{1}{2}} \sum_{n\le y} \frac{\big(\Delta d_2(n;a_1,q_1,a_2,q_2)\big)^2}{n^{\frac{3}{2}}},$$

$$S_{12}(x) = x^{\frac{1}{2}} \sum_{n_1\neq n_2\le y} \frac{\Delta d_2(n_1;a_1,q_1,a_2,q_2)\Delta d_2(n_2;a_1,q_1,a_2,q_2)}{(n_1n_2)^{\frac{3}{4}}} \times \cos\big(4\pi\sqrt{x}(\sqrt{n_1}-\sqrt{n_2})\big).$$

Then

$$\begin{aligned}\int_{\frac{T}{2}}^{T} S_{11}(x)dx &= \sum_{n\le y} \frac{\big(\Delta d_2(n;a_1,q_1,a_2,q_2)\big)^2}{n^{\frac{3}{2}}} \int_{\frac{T}{2}}^{T} x^{\frac{1}{2}}dx \qquad (6.15)\\ &= \sum_{n=1}^{\infty} \frac{\big(\Delta d_2(n;a_1,q_1,a_2,q_2)\big)^2}{n^{\frac{3}{2}}} \int_{\frac{T}{2}}^{T} x^{\frac{1}{2}}dx + O\Big(T^{\frac{3}{2}} \sum_{n>y} \frac{d^2(n)}{n^{\frac{3}{2}}}\Big)\\ &= B_2\big(\frac{a_1}{q_1},\frac{a_2}{q_2}\big) \int_{\frac{T}{2}}^{T} x^{\frac{1}{2}}dx + O(T^{\frac{3}{2}}y^{-\frac{1}{2}}\log^3 y),\end{aligned}$$

noting that $\sum_{n\le N} d^2(n) \ll N\log^3 N$. By using Lemma 3.6 and (6.12), we get

$$\begin{aligned}\int_{\frac{T}{2}}^{T} S_{12}(x)dx &\ll T \sum_{n_1\neq n_2\le y} \frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac{3}{4}}|\sqrt{n_1}-\sqrt{n_2}|} \qquad (6.16)\\ &\ll T\sum_{n\le y} \frac{1}{\sqrt{n+1}-\sqrt{n}} \cdot \frac{d^2(n)}{n^{\frac{3}{2}}}\\ &\ll T\sum_{n\le y} \frac{d^2(n)}{n} \ll T\log^4 y.\end{aligned}$$

From (6.14)-(6.16), we see

$$\int_{\frac{T}{2}}^{T} S_1(x)dx = B_2(\frac{a_1}{q_1},\frac{a_2}{q_2}) \int_{\frac{T}{2}}^{T} x^{\frac{1}{2}}dx + O(T^{\frac{3}{2}}y^{-\frac{1}{2}}\log^3 y), \qquad (6.17)$$

in view of $y \le T\mathcal{L}^{-4}$. Combing (6.10), (6.13) and (6.17), we get

**Lemma 6.2.**

$$\int_{\frac{T}{2}}^{T} R_0^2 dx = \frac{(q_1q_2)^2}{2^6\pi^2} B_2\big(\frac{a_1}{q_1},\frac{a_2}{q_2}\big) \int_{\frac{T}{2}}^{T} x^{\frac{1}{2}}dx + O\big((q_1q_2)^2T^{\frac{3}{2}}y^{-\frac{1}{2}}\log^3 y\big).$$

*6.4. Evaluation of the integral* $\int_{\frac{T}{2}}^{T} R_0(R_1+R_2)dx$

Denote $N = 2^{J+1}H^2$. From (6.7) and (6.8), we see

$$R_0R_1 = \frac{(q_1q_2)^2x^{\frac{1}{2}}}{2^5\pi^2} \sum_{y<n_1\le N} \sum_{n_2\le y} \frac{\cos\big(4\pi\sqrt{n_1x}-\frac{3\pi}{4}\big)\cos\big(4\pi\sqrt{n_2x}-\frac{3\pi}{4}\big)}{(n_1n_2)^{\frac{3}{4}}} \times \Delta d_{2,1}(n_1,H,J;a_1,q_1,a_2,q_2)\Delta d_2(n_2;a_1,q_1,a_2,q_2). \qquad (6.18)$$

Since

$$\begin{aligned}&2\cos\big(4\pi\sqrt{n_1x}-\frac{3\pi}{4}\big)\cos\big(4\pi\sqrt{n_2x}-\frac{3\pi}{4}\big)\\&=\cos\big(4\pi(\sqrt{n_1}-\sqrt{n_2})\sqrt{x}\big)-\sin\big(4\pi(\sqrt{n_1}+\sqrt{n_2})\sqrt{x}\big),\end{aligned}$$

by using (6.12), we get

$$\int_{\frac{T}{2}}^{T}x^{\frac{1}{2}}\cos\big(4\pi\sqrt{n_1x}-\frac{\pi}{4}\big)\cos\big(4\pi\sqrt{n_2x}-\frac{\pi}{4}\big)dx\ll\frac{T}{\sqrt{n_1}-\sqrt{n_2}}.$$

in view of $n_1\neq n_2$. Thus, we have

$$\int_{\frac{T}{2}}^{T}R_0R_1dx\ll(q_1q_2)^2T\sum_{y<n_1\leq N}\sum_{n_2\leq y}\frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac{3}{4}}(\sqrt{n_1}-\sqrt{n_2})}.$$

Dividing the interval $y<n_1\leq N$ into $y<n_1\leq 4y$ and $4y<n_1\leq N$, we see

$$\int_{\frac{T}{2}}^{T}R_0R_1dx\ll(q_1q_2)^2T\Big(\sum\nolimits_5+\sum\nolimits_6\Big),\tag{6.19}$$

where

$$\begin{aligned}\sum\nolimits_5&=\sum_{y<n_1\leq 4y}\sum_{n_2\leq y}\frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac{3}{4}}(\sqrt{n_1}-\sqrt{n_2})},\\\sum\nolimits_6&=\sum_{4y<n_1\leq N}\sum_{n_2\leq y}\frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac{3}{4}}(\sqrt{n_1}-\sqrt{n_2})}.\end{aligned}$$

By the same agrement with (6.16), we get

$$\sum\nolimits_5\ \ll\sum_{\substack{n_1,n_2\leq 4y\\n_1\neq n_2}}\frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac{3}{4}}(\sqrt{n_1}-\sqrt{n_2})}\ll T\log^4y.\tag{6.20}$$

It is clear that

$$\sum\nolimits_6\ll\sum_{4y<n_1\leq N}\frac{d(n_1)}{n_1^{\frac{5}{4}}}\sum_{n_2\leq y}\frac{d(n_2)}{n_2^{\frac{3}{4}}}\ll\log^2y.\tag{6.21}$$

Combining (6.19)-(6.21), we have

$$\int_{\frac{T}{2}}^{T}R_0R_1dx\ll(q_1q_2)^2T\log^4y.$$

The same estimate holds for $\int_{\frac{T}{2}}^{T}R_0R_2dx$, which yields

**Lemma 6.3.**

$$\int_{\frac{T}{2}}^{T}R_0(R_1+R_2)dx\ll(q_1q_2)^2T\log^4y.$$

*6.5. Mean-square of $R_1+R_2$*

From the expression of $R_1$ given by (6.8), we get

$$R_1 \ll x^{\frac{1}{4}}q_1q_2\Big|\sum_{y<n\le N}\frac{e(2\sqrt{nx}-\frac{3}{8})}{n^{\frac{3}{4}}}\Delta d_{2,1}(n,H,J;a_1,q_1,a_2,q_2)\Big|.$$

So we have

$$\begin{aligned}R_1^2 \ll& x^{\frac{1}{2}}(q_1q_2)^2\Big|\sum_{y<n\le N}\frac{e(2\sqrt{nx}-\frac{3}{8})}{n^{\frac{3}{4}}}\Delta d_{2,1}(n,H,J;a_1,q_1,a_2,q_2)\Big|^2\\ =&x^{\frac{1}{2}}(q_1q_2)^2\sum_{y<n_1,n_2\le N}\frac{e\big(2(\sqrt{n_2}-\sqrt{n_1})\sqrt{x}\big)}{(n_1n_2)^{\frac{3}{4}}}\prod_{i=1}^{2}\Delta d_{2,1}(n_i,H,J;a_1,q_1,a_2,q_2).\end{aligned}$$

Splitting the sum according to $n_1=n_2$ or $n_1\neq n_2$, we obtain

$$\begin{aligned}R_1^2 \ll x^{\frac{1}{2}}(q_1q_2)^2\Bigg\{&\sum_{y<n\le N}\frac{d^2(n)}{n^{\frac{3}{2}}}+\sum_{y<n_1<n_2\le N}\frac{e\big(2(\sqrt{n_2}-\sqrt{n_1})\sqrt{x}\big)}{(n_1n_2)^{\frac{3}{4}}}\\ &\times\prod_{i=1}^{2}d_{2,1}(n_i,H,J;a_1,q_1,a_2,q_2)\Bigg\}.\end{aligned}$$

Similarly to (6.12), we get the estimate

$$\int_{\frac{T}{2}}^{T}x^{\frac{1}{2}}e\big(2(\sqrt{n_2}-\sqrt{n_1})\sqrt{x}\big)dx\ll\frac{T}{\sqrt{n_2}-\sqrt{n_1}},$$

which yields

$$\begin{aligned}\int_{\frac{T}{2}}^{T}R_1^2dx \ll&(q_1q_2)^2\sum_{y<n\le N}\frac{d^2(n)}{n^{\frac{3}{2}}}\int_{\frac{T}{2}}^{T}x^{\frac{1}{2}}dx+(q_1q_2)^2\\ &\times\sum_{y<n_1<n_2\le N}\frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac{3}{4}}}\int_{\frac{T}{2}}^{T}x^{\frac{1}{2}}e\big(2(\sqrt{n_2}-\sqrt{n_1})\sqrt{x}\big)dx\\ \ll&T^{\frac{3}{2}}(q_1q_2)^2y^{-\frac{1}{2}}\log^3y+T(q_1q_2)^2\sum_{y<n_1<n_2\le N}\frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac{3}{4}}(\sqrt{n_2}-\sqrt{n_1})},\end{aligned}\tag{6.22}$$

where we used the estimate $\sum\limits_{n\le u}d^2(n)\ll u\log^3u$. By Lemma 3.6, we have

$$\sum_{y<n_1<n_2\le N}\frac{d(n_1)d(n_2)}{(n_1n_2)^{\frac{3}{4}}(\sqrt{n_2}-\sqrt{n_1})}\ll\sum_{y<n\le N}\frac{d^2(n)}{n}\ll\mathcal{L}^4.$$

From (6.22), we obtain the estimate

$$\int_{\frac{T}{2}}^{T}R_1^2dx\ll(q_1q_2)^2T^{\frac{3}{2}}y^{-\frac{1}{2}}\mathcal{L}^3+(q_1q_2)^2T\mathcal{L}^4.$$

For $\int_{\frac{T}{2}}^{T}R_2^2dx$, the same estimate holds. Thus, we get

**Lemma 6.4.**

$$\int_{\frac{T}{2}}^{T} |R_1+R_2|^2 dx \ll (q_1q_2)^2 T^{\frac{3}{2}} y^{-\frac{1}{2}} \mathcal{L}^3 + (q_1q_2)^2 T\mathcal{L}^4.$$

*6.6. Evaluation of the integral* $\int_{\frac{T}{2}}^{T} R^2 dx$

Since $R = R_0 + R_1 + R_2$, we have

$$\int_{\frac{T}{2}}^{T} R^2 dx = \int_{\frac{T}{2}}^{T} R_0^2 dx + 2\int_{\frac{T}{2}}^{T} R_0(R_1+R_2)dx + \int_{\frac{T}{2}}^{T} (R_1+R_2)^2 dx.$$

Then from Lemma 6.2, Lemma 6.3 and Lemma 6.4, it follows that

$$\int_{\frac{T}{2}}^{T} R^2 dx = \frac{(q_1q_2)^2}{2^6\pi^2} B_2\big(\frac{a_1}{q_1}, \frac{a_2}{q_2}\big) \int_{\frac{T}{2}}^{T} x^{\frac{1}{2}} dx + O\big((q_1q_2)^2 T^{\frac{3}{2}} y^{-\frac{1}{2}} \mathcal{L}^3\big).$$

Take $y = T^{\frac{1}{2}}$. We see

**Lemma 6.5.**

$$\int_{\frac{T}{2}}^{T} R^2 dx = \frac{(q_1q_2)^2}{2^6\pi^2} B_2\big(\frac{a_1}{q_1}, \frac{a_2}{q_2}\big) \int_{\frac{T}{2}}^{T} x^{\frac{1}{2}} dx + O\big((q_1q_2)^2 T^{\frac{5}{4}} \mathcal{L}^3\big).$$

*6.7. Mean-value of* $S(q_1q_2x, \frac{a_1}{q_1}, \frac{a_2}{q_2})$

From Lemma 6.5 and Lemma 6.1, using Cauchy-Schwarz inequality, we get

$$\int_{\frac{T}{2}}^{T} |R|G dx \ll (q_1q_2)^2 T^{\frac{5}{4}} \mathcal{L}^2,$$

$$q_1q_2\mathcal{L}^3 \int_{\frac{T}{2}}^{T} |R| dx \ll (q_1q_2)^2 T^{\frac{5}{4}} \mathcal{L}^3,$$

$$\int_{\frac{T}{2}}^{T} (q_1q_2)^2 \mathcal{L}^6 dx \ll (q_1q_2)^2 T\mathcal{L}^6.$$

Combining these three estimates with Lemma 6.5, Lemma 6.1, we obtain

$$\int_{\frac{T}{2}}^{T} S^2(q_1q_2x, \frac{a_1}{q_1}, \frac{a_2}{q_2}) dx = \frac{(q_1q_2)^2}{2^6\pi^2} B_2\big(\frac{a_1}{q_1}, \frac{a_2}{q_2}\big) \int_{\frac{T}{2}}^{T} x^{\frac{1}{2}} dx + O\big((q_1q_2)^2 T^{\frac{5}{4}} \mathcal{L}^3\big),$$

which immediatly implies Theorem 1.4 by noting that the interval $[1, T]$ can be divided into subintervals of the the form $[2^{-j}T, 2^{1-j}T]$, $1 \le j \ll \log T$.

## 7. Proofs of Theorem 1.5

In this section, we shall refer to a method originated from Zhai [17] to prove Theorem 1.5. Throughout this section, we assume $k$ is an integer with $3 \le k < A_0$, where $A_0 > 9$ satisfies (1.5).

### *7.1. Evaluation of the integral* $\int_{\frac{T}{2}}^{T} R_0^k(x;y)dx$

In this subsection, We will evaluate the integral $\int_{\frac{T}{2}}^{T} R_0^k(x;y)dx$, which provides the main term in the asymptotic formula of $\int_{\frac{T}{2}}^{T} S^k(x, \frac{a_1}{q_1}, \frac{a_2}{q_2})dx$.

By the elementary formula

$$(\cos a_1)(\cos a_2)\cdots(\cos a_k) = \frac{1}{2^{k-1}} \sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \cos\big(a_1 + (-1)^{i_1}a_2 + \cdots + (-1)^{i_{k-1}}a_k\big), \tag{7.1}$$

from the expression of $R_0$ shown by (6.7), we get

$$R_0^k(x;y) = \Big(\frac{x^{\frac{1}{4}}q_1q_2}{4\sqrt{2}\pi}\Big)^k \frac{1}{2^{k-1}} \sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \times \sum_{n_i\le y, 1\le i\le k} \frac{\cos\big(4\pi\alpha(\boldsymbol{n};\boldsymbol{i})\sqrt{x} - \beta(\boldsymbol{i})\big)\prod_{j=1}^{k}\Delta d_2(n_j;a_1,q_1,a_2,q_2)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}.$$

Denote

$$S_5(x) = x^{\frac{k}{4}} \sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \sum_{\substack{n_i\le y\\ 1\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})=0}} \frac{\cos\big(\beta(\boldsymbol{i})\big)\prod_{j=1}^{k}\Delta d_2(n_j;a_1,q_1,a_2,q_2)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}, \tag{7.2}$$

$$S_6(x) = x^{\frac{k}{4}} \sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \sum_{\substack{n_i\le y\\ 1\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})\neq 0}} \frac{\cos\big(4\pi\alpha(\boldsymbol{n};\boldsymbol{i})\sqrt{x} - \beta(\boldsymbol{i})\big)\prod_{j=1}^{k}\Delta d_2(n_j;a_1,q_1,a_2,q_2)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}. \tag{7.3}$$

Then $R_0^k(x;y)$ can be rewritten as

$$R_0^k(x;y) = \frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k}\big(S_5(x) + S_6(x)\big). \tag{7.4}$$

First we consider $\int_{\frac{T}{2}}^{T} S_5(x)dx$. By Lemma 3.5, we have

$$\int_{\frac{T}{2}}^{T} S_5(x)dx = \sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \sum_{\substack{n_i\le y\\ 1\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})=0}} \frac{\cos\beta(\boldsymbol{i})}{(n_1n_2\cdots n_k)^{\frac{3}{4}}} \prod_{j=1}^{k}\Delta d_2(n_j;a_1,q_1,a_2,q_2) \int_{\frac{T}{2}}^{T} x^{\frac{k}{4}}dx$$

$$= \sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \sum_{\alpha(\boldsymbol{n};\boldsymbol{i})=0} \cos\beta(\boldsymbol{i}) \prod_{j=1}^{k} \frac{\Delta d_2(n_j;a_1,q_1,a_2,q_2)}{n_j^{\frac{3}{4}}} \int_{\frac{T}{2}}^{T} x^{\frac{k}{4}}dx + O(T^{1+\frac{k}{4}}y^{-\frac{1}{2}+\varepsilon}).$$

Note that

$$\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\sum_{\alpha(\boldsymbol{n};\boldsymbol{i})=0}\cos\beta(\boldsymbol{i})\prod_{j=1}^{k}\frac{\Delta d_2(n_j;a_1,q_1,a_2,q_2)}{n_j^{\frac{3}{4}}}$$

$$=\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\cos\big((k-2|\boldsymbol{i}|)\frac{3\pi}{4}\big)\sum_{\alpha(\boldsymbol{n};\boldsymbol{i})=0}\prod_{j=1}^{k}\frac{\Delta d_2(n_j;a_1,q_1,a_2,q_2)}{n_j^{\frac{3}{4}}}$$

$$=\sum_{\upsilon=1}^{k-1}\cos\big((k-2\upsilon)\frac{3\pi}{4}\big)\binom{k-1}{\upsilon}\sum_{\sqrt{n_1}+\cdots+\sqrt{n_\upsilon}=\sqrt{n_{\upsilon+1}}+\cdots+\sqrt{n_k}}\prod_{j=1}^{k}\frac{\Delta d_2(n_j;a_1,q_1,a_2,q_2)}{n_j^{\frac{3}{4}}}.$$

According to the definition in Section 3, this is $B_k\big(\frac{a_1}{q_1},\frac{a_2}{q_2}\big)$. Thus

$$\int_{\frac{T}{2}}^{T}S_5(x)dx=B_k\big(\frac{a_1}{q_1},\frac{a_2}{q_2}\big)\int_{\frac{T}{2}}^{T}x^{\frac{k}{4}}dx+O(T^{1+\frac{k}{4}}y^{-\frac{1}{2}+\varepsilon}).\tag{7.5}$$

We now consider $\int_{\frac{T}{2}}^{T}S_6(x)dx$. From (7.3), using (6.12), we see

$$\int_{\frac{T}{2}}^{T}S_6(x)dx\tag{7.6}$$

$$=\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\sum_{\substack{n_i\le y,1\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})\neq0}}\prod_{j=1}^{k}\frac{\Delta d_2(n_j;a_1,q_1,a_2,q_2)}{n_j^{\frac{3}{4}}}\int_{\frac{T}{2}}^{T}x^{\frac{k}{4}}\cos\big(4\pi\alpha(\boldsymbol{n};\boldsymbol{i})\sqrt{x}-\beta(\boldsymbol{i})\big)dx$$

$$\ll T^{\frac{1}{2}+\frac{k}{4}}\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\sum_{\substack{n_i\le y,1\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})\neq0}}\frac{d(n_1)\cdots d(n_k)}{(n_1\cdots n_k)^{\frac{3}{4}}|\alpha(\boldsymbol{n};\boldsymbol{i})|}.$$

By Lemma 3.4, we have the estimate

$$\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\sum_{\substack{n_i\le y,1\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})\neq0}}\frac{d(n_1)\cdots d(n_k)}{(n_1\cdots n_k)^{\frac{3}{4}}\alpha(\boldsymbol{n};\boldsymbol{i})}\ll y^{s(k)+\varepsilon}.$$

Substituting it into (7.6), we get

$$\int_{\frac{T}{2}}^{T}S_6(x)dx\ll T^{\frac{1}{2}+\frac{k}{4}}y^{s(k)+\varepsilon}.\tag{7.7}$$

Combining this with (7.4), (7.5), we arrive at

**Lemma 7.1.**

$$\int_{\frac{T}{2}}^{T}R_0^k dx=\frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k}B_k\big(\frac{a_1}{q_1},\frac{a_2}{q_2}\big)\int_{\frac{T}{2}}^{T}x^{\frac{k}{4}}dx+O\Big((q_1q_2)^kT^{\frac{1}{2}+\frac{k}{4}}\big(T^{\frac{1}{2}}y^{-\frac{1}{2}+\varepsilon}+y^{s(k)+\varepsilon}\big)\Big).$$

*7.2. Evaluation of the integral* $\int_{\frac{T}{2}}^{T} R_0^{k-1}(R_1+R_2)dx$

Let $N=2^{J+1}H^2$. From the expressions of $R_0$ and $R_1$ given by (6.7) and (6.8), using (7.1), we have

$$\begin{aligned}R_0^{k-1}R_1 =& \big(\frac{q_1q_2x^{\frac{1}{4}}}{4\sqrt{2}\pi}\big)^k \sum_{y<n_1\le N}\sum_{\substack{n_i\le y\\2\le i\le k}} \frac{\cos(4\pi\sqrt{n_1x}-\frac{3\pi}{4})\cdots\cos(4\pi\sqrt{n_kx}-\frac{3\pi}{4})}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}\\ &\times\Delta d_{2,1}(n_1,H,J;a_1,q_1,a_2,q_2)\prod_{i=2}^{k}\Delta d_2(n_i;a_1,q_1,a_2,q_2)\\ =&\big(\frac{q_1q_2x^{\frac{1}{4}}}{4\sqrt{2}\pi}\big)^k\frac{1}{2^{k-1}}\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\sum_{y<n_1\le N}\sum_{\substack{n_i\le y\\2\le i\le k}}\frac{\cos\big(4\pi\alpha(\boldsymbol{n};\boldsymbol{i})\sqrt{x}-\beta(\boldsymbol{i})\big)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}\\ &\times\Delta d_{2,1}(n_1,H,J;a_1,q_1,a_2,q_2)\prod_{i=2}^{k}\Delta d_2(n_i;a_1,q_1,a_2,q_2).\end{aligned}$$

Set

$$\begin{aligned}S_3(x)=&x^{\frac{k}{4}}\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\sum_{y<n_1\le N}\sum_{\substack{n_i\le y\\2\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})=0}}\frac{\cos\big(\beta(\boldsymbol{i})\big)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}\\ &\times\Delta d_{2,1}(n_1,H,J;a_1,q_1,a_2,q_2)\prod_{i=2}^{k}\Delta d_2(n_i;a_1,q_1,a_2,q_2),\\ S_4(x)=&x^{\frac{k}{4}}\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\sum_{y<n_1\le N}\sum_{\substack{n_i\le y\\2\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})\neq 0}}\frac{\cos\big(4\pi\alpha(\boldsymbol{n};\boldsymbol{i})\sqrt{x}-\beta(\boldsymbol{i})\big)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}}\\ &\times\Delta d_{2,1}(n_1,H,J;a_1,q_1,a_2,q_2)\prod_{i=2}^{k}\Delta d_2(n_i;a_1,q_1,a_2,q_2).\end{aligned}$$

Then

$$R_0^{k-1}R_1=\frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k}\big(S_3(x)+S_4(x)\big). \tag{7.8}$$

From Lemma 3.5, it is easy to get

$$\begin{aligned}\int_{\frac{T}{2}}^{T}S_3(x)dx\ll&\sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}}\sum_{y<n_1\le N}\sum_{\substack{n_i\le y\\2\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})=0}}\prod_{j=1}^{k}\frac{d(n_j)}{n_j^{\frac{3}{4}}}\int_{\frac{T}{2}}^{T}x^{\frac{k}{4}}dx \qquad (7.9)\\ \ll& T^{1+\frac{k}{4}}\sum_{\upsilon=1}^{k-1}\big|s_{k,\upsilon}(d)-s_{k,\upsilon}(d;y)\big|\ll T^{1+\frac{k}{4}}y^{-\frac{1}{2}+\varepsilon}.\end{aligned}$$

We now evaluate $\int_{\frac{T}{2}}^{T} S_4(x)dx$. From (6.12), we have

$$\int_{\frac{T}{2}}^{T} S_4(x)dx \ll T^{\frac{1}{2}+\frac{k}{4}}\Big(\sum\nolimits_1 + \sum\nolimits_2\Big), \tag{7.10}$$

where

$$\sum\nolimits_1 = \sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \sum_{y<n_1\le k^2 y} \sum_{\substack{n_i\le y\\ 2\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})\neq 0}} \frac{d(n_1)\cdots d(n_k)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}|\alpha(\boldsymbol{n};\boldsymbol{i})|},$$

$$\sum\nolimits_2 = \sum_{\boldsymbol{i}\in\mathbf{1}^{k-1}} \sum_{k^2 y<n_1\le N} \sum_{\substack{n_i\le y\\ 2\le i\le k\\ \alpha(\boldsymbol{n};\boldsymbol{i})\neq 0}} \frac{d(n_1)\cdots d(n_k)}{(n_1n_2\cdots n_k)^{\frac{3}{4}}|\alpha(\boldsymbol{n};\boldsymbol{i})|}.$$

By Lemma 3.4, we see

$$\sum\nolimits_1 \ll y^{s(k)+\varepsilon}. \tag{7.11}$$

For $n_1 > k^2 y$, obviously $|\alpha(\boldsymbol{n};\boldsymbol{i})| \gg n_1^{\frac{1}{2}}$ holds. Thus,

$$\sum\nolimits_2 \ll \sum_{k^2 y<n_1\le N} \sum_{\substack{n_i\le y\\ 2\le i\le k}} \frac{d(n_1)\cdots d(n_k)}{(n_2\cdots n_k)^{\frac{3}{4}} n_1^{\frac{5}{4}}} \ll y^{\frac{k-2}{4}}\mathcal{L}^k. \tag{7.12}$$

For $k\ge 3$, we see $s(k) > \frac{k-2}{4}$. Hence, by (7.10)-(7.12), we get

$$\int_{\frac{T}{2}}^{T} S_4(x)dx \ll T^{\frac{1}{2}+\frac{k}{4}} y^{s(k)+\varepsilon}. \tag{7.13}$$

From (7.8), (7.9) and (7.13), it follows that

$$\int_{\frac{T}{2}}^{T} R_0^{k-1}R_1 dx \ll \frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k}\big(T^{1+\frac{k}{4}}y^{-\frac{1}{2}+\varepsilon} + T^{\frac{1}{2}+\frac{k}{4}}y^{s(k)+\varepsilon}\big).$$

Replacing $R_1$ by $R_2$, this estimate still holds, which immediately implies

**Lemma 7.2.**

$$\int_{\frac{T}{2}}^{T} R_0^{k-1}(R_1+R_2)dx \ll (q_1q_2)^k\big(T^{1+\frac{k}{4}}y^{-\frac{1}{2}+\varepsilon} + T^{\frac{1}{2}+\frac{k}{4}}y^{s(k)+\varepsilon}\big).$$

*7.3. Moments of $R_1+R_2$*

Note $K_0\ge A_0$ is an even number. Let $y^{2s(K_0)}\le T$. Then from Lemma 7.1, we see

$$\int_{\frac{T}{2}}^{T} |R_0|^{K_0}dx \ll (q_1q_2)^{K_0}T^{1+\frac{K_0}{4}+\varepsilon}.$$

By Lemma 6.2, we get

$$\int_{\frac{T}{2}}^{T} |R_0|^2 dx \ll (q_1q_2)^2 T^{\frac{3}{2}+\varepsilon}.$$

Using Hölder's inequality, we have

$$\int_{\frac{T}{2}}^{T} |R_0|^{A_0} dx \ll (q_1q_2)^{A_0} T^{1+\frac{A_0}{4}+\varepsilon}. \tag{7.14}$$

Owing to (1.5) and (5.3), we see

$$\int_{\frac{T}{2}}^{T} \big|S\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big)\big|^{A_0} dx \ll (q_1q_2)^{A_0} T^{1+\frac{A_0}{4}+\varepsilon}. \tag{7.15}$$

Thus, from Lemma 6.1, (6.3), (7.14) and(7.15), we have

$$\begin{aligned}\int_{\frac{T}{2}}^{T} |R_1+R_2|^{A_0} dx \ll & \int_{\frac{T}{2}}^{T} \big|S\big(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2}\big)\big|^{A_0} dx + \int_{\frac{T}{2}}^{T} |R_0|^{A_0} dx \\ & + \int_{\frac{T}{2}}^{T} G^{A_0}(x) dx + O\big(T(q_1q_2)^{A_0}\mathcal{L}^{3A_0}\big) \\ \ll & (q_1q_2)^{A_0} T^{1+\frac{A_0}{4}+\varepsilon}.\end{aligned} \tag{7.16}$$

By using Lemma 6.4 and (7.16), and Hölder's inequality,, we obtain

**Lemma 7.3.** *For $T^{\varepsilon} \le y \le T^{\frac{1}{2s(K_0)}}$, we have*

$$\int_{\frac{T}{2}}^{T} |R_1+R_2|^{k} dx \ll (q_1q_2)^{k} T^{1+\frac{k}{4}+\varepsilon} y^{-\frac{A_0-k}{2(A_0-2)}}.$$

*7.4. Evaluation of the integral $\int_{\frac{T}{2}}^{T} R^k dx$*

Since $(a+b)^k = a^k + ka^{k-1}b + O(|a|^{k-2}|b|^2 + |b|^k)$, we have

$$\begin{aligned}\int_{\frac{T}{2}}^{T} R^k dx = & \int_{\frac{T}{2}}^{T} R_0^k dx + k\int_{\frac{T}{2}}^{T} R_0^{k-1}(R_1+R_2) dx \\ & + O\Big(\int_{\frac{T}{2}}^{T} \big|R_0^{k-2}(R_1+R_2)^2\big| dx + \int_{\frac{T}{2}}^{T} \big|R_1+R_2\big|^k dx\Big).\end{aligned}$$

From (7.14), Lemma 7.3 and Hölder's inequality, we get

$$\int_{\frac{T}{2}}^{T} \big|R_0^{k-2}(R_1+R_2)^2\big| dx \ll (q_1q_2)^{k} T^{1+\frac{k}{4}+\varepsilon} y^{-\frac{A_0-k}{2(A_0-2)}}.$$

Combining this two formulas with Lemma 7.1, Lemma 7.2 and Lemma 7.3, we obtain

$$\int_{\frac{T}{2}}^{T} R^k dx = \frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k} B_k(\frac{a_1}{q_1}, \frac{a_2}{q_2}) \int_{\frac{T}{2}}^{T} x^{\frac{k}{4}} dx + O\big((q_1q_2)^{3k} T^{1+\frac{k}{4}+\varepsilon} y^{-\frac{A_0-k}{2(A_0-2)}}\big) + O\big((q_1q_2)^k T^{1+\frac{k}{4}} y^{-\frac{1}{2}+\varepsilon} + (q_1q_2)^k T^{\frac{1}{2}+\frac{k}{4}} y^{s(k)+\varepsilon}\big).$$

Take $y = T^{\frac{1}{2s(K_0)}}$, then $y^{s(k)+\frac{1}{2}} \ll y^{s(K_0)} = T^{\frac{1}{2}}$ and thereby $y^{s(k)} \ll T^{\frac{1}{2}} y^{-\frac{1}{2}}$. Thus, we have the asymptotic formula of $\int_{\frac{T}{2}}^{T} R^k dx$ as following

**Lemma 7.4.**

$$\int_{\frac{T}{2}}^{T} R^k dx = \frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k} B_k(\frac{a_1}{q_1}, \frac{a_2}{q_2}) \int_{\frac{T}{2}}^{T} x^{\frac{k}{4}} dx + O\big((q_1q_2)^k T^{1+\frac{k}{4}-\frac{A_0-k}{4(A_0-2)s(K_0)}+\varepsilon}\big).$$

*7.5. Moments of* $S(q_1q_2x; \frac{a_1}{q_1}, \frac{a_2}{q_2})$

From (7.14) and Lemma 6.1, by Hölder's inequality, we get

$$\int_{\frac{T}{2}}^{T} |R|^{k-1} G dx \ll (q_1q_2)^k T^{\frac{3}{4}+\frac{k}{4}+\varepsilon}, \tag{7.17}$$

$$q_1q_2\mathcal{L}^3 \int_{\frac{T}{2}}^{T} |R|^{k-1} dx \ll (q_1q_2)^k T^{\frac{3}{4}+\frac{k}{4}+\varepsilon}. \tag{7.18}$$

Note that

$$\int_{\frac{T}{2}}^{T} (q_1q_2)^k \mathcal{L}^{3k} dx \ll (q_1q_2)^k T^{1+\varepsilon}. \tag{7.19}$$

Since $\frac{A_0-k}{4(A_0-2)s(K_0)} < \frac{1}{4}$ for $3 \le k < A_0$, from (6.3), Lemma 7.4, Lemma 6.1, (7.17)-(7.19), we can conclude that

$$\int_{\frac{T}{2}}^{T} S^k(q_1q_2x, \frac{a_1}{q_1}, \frac{a_2}{q_2}) dx = \frac{(q_1q_2)^k}{2^{\frac{7}{2}k-1}\pi^k} B_k(\frac{a_1}{q_1}, \frac{a_2}{q_2}) \int_{\frac{T}{2}}^{T} x^{\frac{k}{4}} dx + O\big((q_1q_2)^k T^{1+\frac{k}{4}-\frac{A_0-k}{4(A_0-2)s(K_0)}+\varepsilon}\big).$$

Then we get the evaluation of the integral $\int_1^T S^k(q_1q_2x, \frac{a_1}{q_1}, \frac{a_2}{q_2}) dx$ by summing over integrals over intervals of the form $[2^{-j}T, 2^{1-j}T]$, $1 \le j \ll \log T$.

## References


[1] B. C. Berndt, S. Kim, A. Zaharescu, Weighted divisor sums and Bessel function series, III, Journal fr die reine und angewandte Mathematik (Crelles Journal) 683 (2013) 67-96.

[2] H. Cramér. Über zwei Sätze des Herrn G. H. Hardy. Mathematische Zeitschrift, 15(1):201-210, 1922.

[3] D. R. Heath-Brown. The distribution and moments of the error term in the dirichlet divisor problem. Acta Arith, 60(4):389-415, 1992.

[4] M. N. Huxley. Exponential sums and lattice points III. Proceedings of the London Mathe- matical Society, 87(03):591-609, 2003.

[5] M. N. Huxley. Exponential sums and the Riemann zeta function V. Proceedings of the London Mathematical Society, 90(01):1-41, 2005.

[6] A. Ivić. Large values of the error term in the divisor problem. Inventiones mathematicae, 71(3):513-520, 1983.

[7] A. Ivic. The Riemann zeta-function: theory and applications. Courier Dover Publications, 2003.

[8] A. Ivi c and P. Sargos. On the higher moments of the error term in the divisor problem. Illinois Journal of Mathematics, 51(2):353-377, 2007.

[9] K. Liu. On higher-power moments of the error term for the divisor problem with congruence conditions. Monatshefte fr Mathematik, 163(2):175-195, 2011.

[10] W. Mller and W. G. Nowak. Third power moments of the error terms corresponding to certain arithmetic functions. manuscripta mathematica, 87(1):459-480, 1995.

[11] H. E. Richert. Ein Gitterpunktproblem. Mathematische Annalen, 125(1):467-471, 1952.

[12] S.H. Min. The methods of number thoery (in Chinese). Science Press, 1981.

[13] Z. Shan. Hilbert inequality (in chinese). Chinese Science Bulletin, 29(01):021, 1984.

[14] K. M. Tsang. Higher-power moments of (x), e(t) and p(x). Proceedings of the London Math- ematical Society, 65:65-84, 1992.

[15] J. D. Vaaler. Some extremal functions in fourier analysis. Bulletin of the American Mathe- matical Society, 12(2):183-216, 1985.

[16] G. Voronoï. Sur une fonction transcendante et ses applications à la sommation de quelques séries. In Annales scientifiques de l'École Normale Supérieure, volume 21, pages 207-267, 1904.

[17] W. G. Zhai. On higher-power moments of $\Delta(x)$(II). Acta Arith, 114:35-54, 2004.

[18] W. G. Zhai. On the error term in weyls law for heisenberg manifolds. Acta Arithmetica, 134:219-257, 2008.